

Author Index

- Abouleish A. Postoperative management of intraoperative anaphylaxis (letter), 92:559
- Abramson DC, see Pivalizza EG
- Acalovschi I, Cristea T, Margarit S, Gavrut R. Tramadol added to lidocaine for intravenous regional anesthesia, 92:209
- Adachi YU
Higuchi H. Continuous infusion of prostaglandin E₁ and epidural analgesia for a patient with paroxysmal nocturnal hemoglobinuria (letter), 92:1618
- Watanabe K, Higuchi H, Satoh T. The determinants of propofol induction of anesthesia dose, 92:656
see Higuchi H
- Adler LJ, see Vallejo MC
- Agashe GS, see Kingery WS
- Ahlén KG, see Finucane BT
- Ahluwalia A, see Taguchi A
- Ahmaidi S, see Ayoub J
- Ahmed HE, see White PF
- Akça O, see Goll V
- Alexander JM, Sharma SK, McIntire DD, Wiley J, Leveno KJ. Intensity of labor pain and cesarean delivery, 92:1524
- Alexander R, see Gan TJ
- Alfery DD, see Seitz PA
- Ali AA, Ali GS, Steinke JM, Shepherd AP. Co-oximetry interference by hemoglobin-based blood substitutes, 92:863
- Ali GS, see Ali AA
- Allaouchiche B, see Debon R; Duflo F
- Allen MG, see Kaushik S
- Allvin R, see Rawal N
- Alonchin A, see Eilig I
- Amann A, see Rieder J
- Amilon A, see Rawal N
- Andersen O, see Bruun LS
- Ando-Nagase K, see Nagase K
- Andropoulos DA, see Stayer SA
- Andropoulos DB, see Stayer SA
- Angermeier KW, see O'Hara JF Jr
- Ang E-T, see Langeron O
- Ankri A, A, see Huraux C
- Anusit J, see Iamaroon A
- Aouad MT, Siddik SS, Jalbout MI, Baraka AS. Does pregnancy protect against intrathecal lidocaine-induced transient neurologic symptoms?, 92:401
- Siddik SS, Rizk LB, Zaytoun GM, Baraka AS. The effect of dexamethasone on postoperative vomiting after tonsillectomy, 92:636
- Apfel CC, see Greim C-A
- Appavu SK, see Khorasani A
- April E, see Vloka JD
- Aram L, Krane EJ, Kozloski LJ, Yaster M. Tunneled epidural catheters for prolonged analgesia in pediatric patients, 92:1432
- Aregawi A, see Patel RI
- Argiriadou H, see Himmelseher S
- Arkiliç CF, see Goll V; Taguchi A
- Arlander E, see Modalen ÅÖ
- Armbruster SC, see Coloma M
- Armstead VE, see Nielsen VG
- Aron E, see Holzman RS
- Artru AA, see Eilig I
- Asada A, see Kariya N
- Asai T, Yamamoto K, Hirose T, Taguchi H, Shingu K. Breakage of epidural catheters: a comparison of an arrow reinforced catheter and other nonreinforced catheters, 92:246
- Atallah F, Bastide-Heulin T, Samii K, Plante P. Nefopam analgesia as a substitute of neurolept analgesia for extracorporeal shock wave lithotripsy (letter), 92:557
- Aubert J-D, see Broccard AF
- Awad AA, Ghobashy AM, Ouda W, Stout RG, Silverman DG, Shelley KH. Different responses of ear and finger pulse oximeter wave form to cold pressor test, 92:1483
- Ayoub J, Cohendy R, Prioux J, Ahmaidi S, Bourgeois JM, Dauzat M, Ramonatxo M, Préfaut C. Diaphragm movement before and after cholecystectomy: a sonographic study, 92:755
- Babin D, see Bernard F
- Bacon DR, see Batt RE
- Bahk J-H, Lim Y-J. Use of intracuff lidocaine during general anesthesia (letter), 92:1075
- Balea MC, see Russell IA
- Bali A, see Helmy SAK
- Baliga N, see Wahr JA
- Bamber J, see Chilvers RJ
- Bannister CF, Brosius KK, Sigl JC, Meyer BJ, Sebel PS. The effect of bispectral index monitoring on anesthetic use and recovery in children anesthetized with sevoflurane in nitrous oxide, 92:877
- Bänziger O, see Mönkhoff M
- Baraka AS, see Aouad MT
- Baranov D. Textbook of Neuroanaesthesia and Critical Care (book review), 92:286
- Barnett RA, see Ochroch EA
- Barthe A, see Bernard J-M
- Barth MC, see Kadar AG
- Bartkowski R, see McNulty SE
- Bartoli M, see Sia S
- Barzilai B, see Skubas NJ
- Barzilay Z, see Vaknin Z
- Basto ER, Waintrop C, Mourey FD, Landru JP, Eurin BG, Jacob LP. Intravenous ketoprofen in thyroid and parathyroid surgery, 92:1052
- Bastide-Heulin T, see Atallah F
- Batt RE, Bacon DR. Clarence J. Durshordwe, the International Anesthesia Research Society, and the World Federation of Societies of Anesthesiologists: the last true disciple of Francis Hoeffler McMechan, 92:1349
- Bauer H, see Fitzgerald RD
- Beardslee M, see Skubas NJ
- Beaussier M, see Coriat P
- Beccaria P, see Casati A
- Becke K, see Schmitt HJ
- Bedforth N, Hardman JG, Nathanson M. Cerebral hemodynamic response to the introduction of desflurane: a comparison with sevoflurane (letter, reply), 92:282
- Behne M, see Bremerich DH
- Bekker AY, Kaufman B, Samir H, Doyle W. The use of dexmedetomidine infusion for awake craniotomy, 92:1251
- Ben-Abraham R, see Vaknin Z
- Benckart D, see Heres EK
- Ben-David B, Stonebraker VC, Hershman R, Frost CL, Williams HK. Survival after failed intraoperative resuscitation: a case of Lazarus syndrome, 92:690
- Bennett-Guerrero E, see Sekaran P
- Bennett J, see Butterworth J
- Benson M, see Junger A
- Bent ST, see Stayer SA
- Berenholtz SM, see Pronovost PJ
- Berkenbosch JW, Lam J, Burd RS, Tobias JD. Noninvasive monitoring of carbon dioxide during mechanical ventilation in older children: end-tidal versus transcutaneous

- techniques, 92:1427
- Berkenstadt H, Margalit N, Hadani M, Friedman Z, Segal E, Villa Y, Perel A. Stroke volume variation as a predictor of fluid responsiveness in patients undergoing brain surgery, 92:984
- Bernard F, Denault A, Babin D, Goyer C, Couture P, Couturier A, Buithieu J. Diastolic dysfunction is predictive of difficult weaning from cardiopulmonary bypass, 92:291
- Bernard J-M, Le Roux D, Barthe A, Jourdain O, Vizquel L, Michel C. The dose-range effects of sufentanil added to 0.125% bupivacaine on the quality of patient-controlled epidural analgesia during labor, 92:184
- Bernstein M, see Blanshard HJ
- Berry A, see Brimacombe J
- Bert A, see Maslow A
- Berti M, see Casati A
- Bertrand M, Godet G, Meerschaert K, Brun L, Salcedo E, Coriat P. Should the angiotensin II antagonists be discontinued before surgery?, 92:26
- Béye-Basse A, see Junger A
- Bhatia S, see Munis JR
- Bhavani-Shankar K, Harnett MJP
- Birbaumer N, see Kiefer R-T
- Birnback DJ, see Browne IM; Schneider MC
- Bitsch-Larsen D, see Bruun LS
- Blackley O, Mychaskiw G II. A convenient way to secure intravenous, arterial, and other lines on extremities (letter), 92:1355
- Blanshard HJ, Chung F, Manninen PH, Taylor MD, Bernstein M. Awake craniotomy for removal of intracranial tumor: considerations for early discharge, 92:89
- Blas ML
- Lobato E. Postoperative management of intraoperative anaphylaxis (letter, reply), 92:559
- see Janelle GM
- Bloos F, see Schwarzkopf K
- Body SC, see Kallmeyer IJ
- Boehler M, see Rieder J
- Boggs DP, see Robins B
- Boldt J, see Haisch G; Lehmann A
- Bonke B, see Kerssens C
- Bonnet F, see Junca A; Langeron O
- Bonnichon P, see Dieudonne N
- Booke M, Van Aken H, Storm M, Fritzsche F, Wirtz S, Hinder F. Fat elimination from autologous blood, 92:341
- Booth JV, see Reynolds JD
- Bordet F, see Duflo F
- Borgeat A, Kalberer F, Jacob H, Ruetsch YA, Gerber C. Patient-controlled interscalene analgesia with ropivacaine 0.2% versus bupivacaine 0.15% after major open shoulder surgery: the effects on hand motor function, 92:218
- Borger MA, see Hynninen M
- Boselli E, see Debon R
- Bourgeois JM, see Ayoub J
- Bovill JG, see White PF
- Bowdle TA, see Cheng DCH; Howie MB
- Bowie RA, O'Connor PJ, Hardman JG, Mahajan RP. The effect of continuous positive airway pressure on cerebral blood flow velocity in awake volunteers, 92:415
- Bradley EL Jr, see Kissin I
- Brannath W, see Gustorff B
- Bräunlich P, see Franz A
- Braun S, see Dietrich W
- Bremerich DH, Neidhart G, Heimann K, Kessler P, Behne M. Prophylactically-administered rectal acetaminophen does not reduce postoperative opioid requirements in infants and small children undergoing elective cleft palate repair, 92:907
- Bremer M, see Fischer LG
- Bresing KA, see Hering R
- Bright CA, see Kissin I
- Brimacombe J, Keller C, Berry A. Gastric insufflation with the ProSeal laryngeal mask, 92:1614
- Broccard AF, Liaudet L, Aubert J-D, Schnyder P, Schaller M-D. Negative pressure post-tracheal extubation alveolar hemorrhage, 92:273
- Brodner G, Van Aken H, Hertle L, Fobker M, Von Eckardstein A, Goeters C, Buerkle H, Harks A, Kehlet H. Multimodal perioperative management—combining thoracic epidural analgesia, forced mobilization, and oral nutrition—reduces hormonal and metabolic stress and improves convalescence after major urologic surgery, 92:1594
- Bromm B, see Vahle-Hinz C
- Brooker RF, see Sherman SV
- Brosius KK, see Bannister CF
- Browne IM, Birnback DJ. Unmasked mischief (letter), 92:279
- Bruguerolle B, Paut O
- Brun L, see Bertrand M
- Bruun LS, Elkjaer S, Bitsch-Larsen D, Andersen O. Hepatic failure in a child after acetaminophen and sevoflurane exposure, 92:1446
- Buerkle H, see Brodner G
- Bufler J, see Haeseler G
- Buithieu J, see Bernard F
- Bukenya D, see Cheng DCH; Howie MB
- Bumpous JM, see Mortero RF
- Burd RS, see Berkenbosch JW
- Burger R, see Simma B
- Burkey DR. Practical Management of Pain, 3rd Edition (book review), 92:561
- Butterworth J
- James RL, Lin YA, Bennett J, Prielipp RC. Gender does not influence epsilon-aminocaproic acid concentrations in adults undergoing cardiopulmonary bypass, 92:1384
- see Groban L; Sherman SV
- Cahalan MK, see Miller JP; Russell IA
- Calderón E, Pernia A, De Antonio P, Calderón-Pla E, Torres L-M. A comparison of two constant-dose continuous infusions of remifentanyl for severe postoperative pain, 92:715
- Calderón-Pla E, see Calderón E
- Camboulives J, Paut O
- Campbell DC. Potencies and probabilities: one-sided *P* values suggest a one-sided story! (letter, reply), 92:278
- Campbell JN, see Shir Y
- Candido KD, see Khorasani A
- Capdevila X, see Langeron O
- Cardigan R, see Royston D
- Carles M, Pulcini A, Macchi P, Duflos P, Raucoules-Aime M, Grimaud D. An evaluation of the brachial plexus block at the humeral canal using a neurostimulator (1417 patients): the efficacy, safety, and predictive criteria of failure, 92:194
- Carli F
- Perioperative Medicine; Volume 18, Number 3 of *Anesthesiology Clinics of North America* (book review), 92:561
- see Finucane BT
- Carli PA, see Legros CB
- Carroll J-A, see Hynninen M
- Casati A, Fanelli G, Magistrali L, Beccaria P, Berti M, Torri G. Minimum local anesthetic volume blocking the femoral nerve in 50% of cases: a double-blinded comparison between 0.5% ropivacaine and 0.5% bupivacaine, 92:205
- Cassola L, see Russell IA
- Casta A, see Gruber EM
- Cathomas G, see Zaugg M

- Cechetto DF, Diab T, Gibson CJ, Gelb AW.
The effects of propofol in the area
postrema of rats, 92:934
- Cespedes RM, see Lathi KG
- Chaiyaroj O, see Iamaron A
- Chakravorti S, see Gruber EM
- Chan C-C, see Tang C-S
- Chaney JD, Nielsen VG. Considerations
for the hemophiliac patient with
inhibitors to factor VIII, 92:785
- Chaney MA, see Chapochnikov AN
- Chang C-F, see Liu Y-H
- Chang C-L, see Chen T-Y
- Chang P-J, see Tsai Y-C
- Chapochnikov AN, Chaney MA.
Transoesophageal
Echocardiography in Anesthesia
(book review), 92:562
- Charlesworth DC, see Surgenor SD
- Chassard D, see Debon R; Duflo F
- Chen B-S, see Chen T-Y
- Cheng CJC, Garewal DS. A failure of the
chain-link mechanism on the
Ohmeda Excel 210 anesthetic
machine, 92:913
- Cheng D, see Howie MB
- Cheng DCH
Newman MF, Duke P, Wong DT,
Finegan B, Howie M, Fitch J,
Bowdle TA, Hogue C, Hillel Z,
Pierce E, Bukenya D. The efficacy
and resource utilization of
remifentanyl and fentanyl in fast-
track coronary artery bypass graft
surgery: a prospective randomized,
double-blinded controlled, multi-
center trial, 92:1094
see Hynninen M
- Chen J, Xu W, Jiang H. 17 β -Estradiol
protects neurons from ischemic
damage and attenuates
accumulation of extracellular
excitatory amino acids, 92:1520
- Chen J-C, see Li J-Y
- Chen L-K, see Yeh H-M
- Chen PP, Chui PT, Ma M, Gin T. A
prospective survey of patients after
cessation of patient-controlled
analgesia, 92:224
- Chen T-Y, Tseng C-C, Wang L-K, Tsai T-Y,
Chen B-S, Chang C-L. The clinical
use of small-dose tetracaine spinal
anesthesia for transurethral
prostatectomy, 92:1020
- Chen Y-L, see Li J-Y
- Chester AH, see Forrest MR
- Chilvers RJ, Bamber J. Postdural puncture
headache in obstetric patients
(letter), 92:1616
- Chiravuri SD, see Nimmagadda U
- Chiu JW, see Coloma M; White PF; Zárate
E
- Choi PT-L, Halpern SH, Malik N, Jadad
AR, Tramèr MR, Walder B.
Examining the evidence in
anesthesia literature: a critical
appraisal of systematic reviews,
92:700
- Choi SS, see Finkel JC
- Christakis GT, see Hynninen M
- Chui PT, see Chen PP
- Chung F, see Blanshard HJ
- Clarke D, see Wadhwa A
- Clark JD, see Kingery WS
- Clark LD, see Mortero RF
- Clough RA, see Surgenor SD
- Coates KM, Mather LE, Johnson R, Flood
P. Thiopental is a competitive
inhibitor at the human $\alpha 7$ nicotinic
acetylcholine receptor, 92:930
- Cobo B, see Fernández-Guisasola J
- Cohendy R, see Ayoub J
- Cohen IT, see Finkel JC
- Cohen S, see Doss NW
- Colburn WA, see O'Hara JF Jr
- Coleman EJ, see Fischer LG
- Collard CD, see Kallmeyer JJ
- Coloma M
Chiu JW, White PF, Armbruster SC. The
use of esmolol as an alternative to
remifentanyl during desflurane
anesthesia for fast-track outpatient
gynecologic laparoscopic surgery,
92:352
- Duffy LL, White PF, Tongier WK, Huber
PJ Jr. Dexmethasone facilitates
discharge after outpatient anorectal
surgery, 92:85
- Columb MO
Polley LS. Potencies and probabilities:
one-sided *P* values suggest a one-
sided story! (letter), 92:278
see Robinson AP
- Comunale ME, see Park KW
- Conrad B, see Fischer LG
- Cook DJ, see Rehfeldt KH
- Cook DR, Dhaliwal DK, Davis PJ, Davis J.
Anesthetic interference with laser
function during excimer laser
procedures in children, 92:1444
- Cooper G, see Maslow A
- Coriat P
Beaussier M. Fast-tracking after
coronary artery bypass graft
surgery, 92:1081
see Bertrand M; David J-S; Huraux C;
Langeron O
- Couture P, see Bernard F
- Couturier A, see Bernard F
- Craig WF, see White PF
- Crawford CC. Awareness during
Anesthesia (book review), 92:1079
- Crimi T, see Doss NW
- Cristea T, see Acalovschi I
- Crystal GJ, see Nimmagadda U
- Cucchiario G, De Lagausie P, El-Ghonemi
A, Nivoche Y. Single-dose caudal
anesthesia for major intraabdominal
operations in high-risk infants,
92:1439
- Culebras X, Van Gessel E, Hoffmeyer P,
Gamulin Z. Clonidine combined
with a long acting local anesthetic
does not prolong postoperative
analgesia after brachial plexus block
but does induce hemodynamic
changes, 92:199
- Dacey LJ, see Surgenor SD
- Dagnino JA, see Muñoz HR
- Dai HB, see Park KW
- Dakin D, see Tweed WA
- Dalibon N, see Liu N
- Daneshgari V, see Zárate E
- Datta S, Harnett MJP
- Dauzat M, see Ayoub J
- David J-S, Vivien B, Lecarpentier Y, Coriat
P, Riou B. Extracellular calcium
modulates the effects of protamine
on rat myocardium, 92:817
- Davies MF, see Kingery WS
- Davis J, see Cook DR
- Davis L
Ross N. Bilateral vocal cord palsy after
ventricular drainage in a child,
92:358
Unlighted stylet tracheal intubation
(letter, reply), 92:1356
- Davis PJ, see Cook DR; Gruber EM
- Deal DD, see Groban L
- De Antonio P, see Calderón E
- Debon R
Allaouchiche B, Duflo F, Boselli E,
Chassard D. The analgesic effect of
sufentanil combined with
ropivacaine 0.2% for labor
analgesia: a comparison of three
sufentanil doses, 92:180
see Duflo F
- Defoe GR, see Surgenor SD
- Degesves S, see Vergnion M
- de Jongh R, see Vranken JH
- De Kock M, see Singelyn FJ
- De Lagausie P, see Cucchiario G
- de L Dear G, see Gan TJ
- de Leon-Casasola OA, see de Médicis E
- Deloof T, see De Witte JL
- de Médicis E, de Leon-Casasola OA.

- Reversible paraplegia associated with lumbar epidural analgesia and thoracic vertebral metastasis, 92:1316
- Demopoulos G, see Grond S
- Denault A, see Bernard F
- Dengler R, see Haeseler G
- Deng X, see Xiao W
- Denson DD, see Hord AH; Kaushik S
- DeSilva A, see Russell IA
- Detsch O, see Vahle-Hinz C
- Deviatko E, see Gruber EM
- De Witte JL, Schoenmaekers B, Sessler DI, Deloof T. The analgesic efficacy of tramadol is impaired by concurrent administration of ondansetron, 92:1319
- Dexter F
- Epstein RH, Marsh HM. A statistical analysis of weekday operating room anesthesia group staffing costs at nine independently managed surgical suites, 92:1493
- Epstein RH, Penning DH. Statistical analysis of postanesthesia care unit staffing at a surgical suite with frequent delays in admission from the operating room—a case study, 92:947
- Gan TJ, Naguib M, Lubarsky DA. Cost identification analysis for succinylcholine, 92:693
- Macario A, Lubarsky DA. The impact on revenue of increasing patient volume at surgical suites with relatively high operating room utilization, 92:1215
- Penning DH, Traub RD. Statistical analysis by Monte-Carlo simulation of the impact of administrative and medical delays in discharge from the postanesthesia care unit on total patient care hours, 92:1222
- Traub RD, Lebowitz P. Scheduling a delay between different surgeons' cases in the same operating room on the same day using upper prediction bounds for case durations, 92:943
- Dhaliwal DK, see Cook DR
- Dhileepan S, Stacey RGW. A preliminary investigation of remifentanyl as a labor analgesic (letter), 92:1358
- Diab T, see Cechetto DF
- Dietrich G, see Junger A
- Dietrich W, Braun S, Spannagl M, Richter JA. Low preoperative antithrombin activity causes reduced response to heparin in adult but not in infant cardiac-surgical patients, 92:66
- Dieudonne N, Gomola A, Bonnichon P, Ozier YM. Prevention of postoperative pain after thyroid surgery: a double-blind randomized study of bilateral superficial cervical plexus blocks, 92:1538
- Dimaculangan DP, Lokhandwala BS, Wlody DJ, Gross RM. Difficult airway in a patient with Coffin-Siris syndrome, 92:554
- Doelberg M, see Langeron O
- Dohi S, see Ma H-C; Yanagidate F
- Dolinski SY, Gerancher JC. Unmasked mischief (letter, reply), 92:280
- Dong Y, see Fiechtner BK
- Dorman T, see Pronovost PJ
- Doss NW, Ipe J, Crimi T, Rajpal S, Cohen S, Fogler RJ, Michael R, Gintautas J. Continuous thoracic epidural anesthesia with 0.2% ropivacaine versus general anesthesia for perioperative management of modified radical mastectomy, 92:1552
- Douvdevani A, see Lewis E
- Doyle W, see Bekker AY
- Driessen B, see Jahr JS
- Duffy LL, see Coloma M
- Duflo F
- Allaouchiche B, Debon R, Bordet F, Chassard D. An evaluation of the gram stain in protected bronchoalveolar lavage fluid for the early diagnosis of ventilator-associated pneumonia, 92:442
- see Debon R
- Duflos P, see Carles M
- Duke P, see Cheng DCH
- Dupont H, see Lang-Lazdunski L
- Dupuis J-Y. Against the discontinuation of angiotensin II antagonists before surgery (letter), 92:1616
- Durcan S, see Swenson JD
- Durieux ME, see Fischer LG; Hollmann MW; Liu H-T
- Dutton R, see Zhang Y
- Dy EA, see Samra SK
- Dzwonczyk R, see Howie MB
- Efrati O, see Vaknin Z
- Eger EI II
- Laster MJ. The effect of rigidity, shape, unsaturation, and length on the anesthetic potency of hydrocarbons, 92:1477
- see Zhang Y
- Ehrlich M, see Sekaran P
- Eilig I, Rachinsky M, Artru AA, Alonchin A, Kapuler V, Tarnapolski A, Shapira Y. The effect of treatment with albumin, hetastarch, or hypertonic saline on neurological status and brain edema in a rat model of closed head trauma combined with uncontrolled hemorrhage and concurrent resuscitation in rats, 92:669
- Eisenhardt S, see Martens-Lobenhoffer J
- Elam M, see Thorén A
- El-Ghoniemi A, see Cucchiari G
- Elia N, see Walder B
- Elkjaer S, see Bruun LS
- Elliott P, Mirakhor RK. Remifentanyl in cardiac surgery (letter, reply), 92:558
- El-Mansouri M, see Reuben SS
- El-Moalem H, see Gan TJ
- El-Orbany MI, see Nimmagadda U
- Engel JM, Hohaus T, Ruwoldt R, Menges T, Jürgensen I, Hempelmann G. Regional hemostatic status and blood requirements after total knee arthroplasty with and without tranexamic acid or aprotinin, 92:775
- Epstein RH, see Dexter F
- Ereth MH, see Fiechtner BK
- Eriksson H, see Laisalmi M
- Eriksson LI, see Modalen ÅÖ
- Erlacher W, see Marhofer P
- Esakof DD, see Lathi KG
- Eurin BG, see Basto ER
- Evans SF, see Pavy TJG
- Eyraud D, see Huraux C
- Fabiani R, see Koessler MJ
- Falk M, see Simma B
- Fanconi S, see Mönkhoff M; Simma B
- Fanelli G, see Casati A
- Fanning GL. Hyaluronidase in ophthalmic anesthesia (letter), 560
- Fan S-Z, see Tang C-S
- Fazi L, Jantzen EC, Rose JB, Kurth CD, Watcha MF. A comparison of oral clonidine and oral midazolam as preanesthetic medications in the pediatric tonsillectomy patient, 92:56
- Felfernig M, see Franz A
- Felleiter P, see Gustorff B
- Feng W, see Maslow A
- Fernández-Guisasola J, Serrano ML, Cobo B, Muñoz L, Plaza A, Trigo C, García del Valle S. A comparison of 0.0625% bupivacaine with fentanyl and 0.1% ropivacaine with fentanyl for continuous epidural labor analgesia, 92:1261
- Fetherston J, see Holzmann RS
- Fiechtner BK, Nuttall GA, Johnson ME, Dong Y, Sujirattanawimol N, Oliver

- WC Jr, Sarpal RS, Oyen LJ, Ereth MH. Plasma tranexamic acid concentrations during cardiopulmonary bypass, 92:1131
- Filipovic M, see Zwetsch G
- Fillingner M, see Surgenor SD
- Finegan B, see Cheng DCH
- Finegold H, see Vallejo MC
- Finkel JC, Cohen IT, Hannallah RS, Patel KM, Kim MS, Hummer KA, Choi SS, Pena M, Schreiber SB, Zalzal G. The effect of intranasal fentanyl on the emergence characteristics after sevoflurane anesthesia in children undergoing surgery for bilateral myringotomy tube placement, 92:1164
- Finucane BT
- Ganapathy S, Carli F, Pridham JN, Ong BY, Shukla RC, Kristofferson AHM, Huizar KM, Nevin K, Ahlén KG, and the Canadian Ropivacaine Research Group. Prolonged epidural infusions of ropivacaine (2 mg/mL) after colonic surgery: the impact of adding fentanyl, 92:1276
- see O'Connor PJ
- Fischer LG
- Bremer M, Coleman EJ, Conrad B, Krumm B, Gross A, Hollmann MW, Mandell G, Durieux ME. Local anesthetics attenuate lysophosphatidic acid-induced priming in human neutrophils, 92:1041
- see Strunk V
- Fischler M, see Liu N
- Fisher D. Statistical clarification (letter), 92:1073
- Fitch J, see Cheng DCH
- Fitzgerald RD, Lamm C, Oczenski W, Stimpfl T, Vycudilik W, Bauer H. Direct current auditory evoked potentials during wakefulness, anesthesia, and emergence from anesthesia, 92:154
- Fitzsimons MG, Peterfreund RA, Raines DE. Aprotinin administration and pulmonary thromboembolism during orthotopic liver transplantation: report of two cases, 92:1418
- Fletcher H, see Myles PS
- Flood P, see Coates KM
- Fobker M, see Brodner G
- Fogler RJ, see Doss NW
- Forrest MR, Chester AH, Royston D. The effects of diaspirin cross-linked hemoglobin on the tone of human saphenous vein, 92:324
- Fox JA, see Kallmeyer JJ
- Franz A, Bräunlich P, Gamsjäger T, Felfernig M, Gustorff B, Kozek-Langenecker SA. The effects of hydroxyethyl starches of varying molecular weights on platelet function, 92:1402
- Fraser CD, see Stayer SA
- Fredman B, Zohar E, Tarabykin A, Shapiro A, Mayo A, Klein E, Jedeikin R. Bupivacaine wound instillation via an electronic patient-controlled analgesia device and a double-catheter system does not decrease postoperative pain or opioid requirements after major abdominal surgery, 92:189
- Freitag H, see Goll V; Marhofer P
- Friedman Z, see Berkenstadt H
- Fritz HG, see Schwarzkopf KRG
- Fritzsche F, see Booke M
- Frost CL, see Ben-David B
- Fujii H, see Yokoyama M
- Fujii T, see Yokota K
- Fujii Y
- Hoshi T, Toyooka H. Colforsin daropate improves contractility in fatigued canine diaphragm, 92:762
- Hoshi T, Uemura A, Toyooka H. Dose-response characteristics of midazolam for reducing diaphragmatic contractility, 92:1590
- Fujioka S, Sasaki T, Hirata K, Okada K, Takazawa A, Sakahashi H, Toyama A. The relationship between plasma concentration of mature adrenomedullin and jugular venous oxygen saturation during and after cardiopulmonary bypass, 92:837
- Fujisawa J, see Inada T
- Fujita S, see Nakae Y
- Fukuda T, Nakayama H, Yanagi K, Mizutani T, Miyabe M, Ohshima N, Toyooka H. The effects of 30% and 60% xenon inhalation on pial vessel diameter and intracranial pressure in rabbits, 92:1245
- Fukusaki M, see Hara T
- Fukushima T, see Tokioka H
- Furuya H, see Yoshitani K
- Gamsjäger T, see Franz A
- Gamulin Z, see Culebras X
- Ganapathy S, see Finucane BT
- Gan TJ
- Madan R, Alexander R, Jhaveri R, El-Moalem H, Weatherwax K, Glass PSA. Duration of action of vecuronium after an intubating dose of rapacuronium, vecuronium, or succinylcholine, 92:1199
- Sloan F, de L Dear G, El-Moalem HE, Lubarsky DA. How much are patients willing to pay to avoid postoperative nausea and vomiting?, 92:393
- see Dexter F
- Garcet L, see Vergnion M
- García del Valle S, see Fernández-Guisasola J
- Gare M, see Gross ER
- Garewal DS, see Cheng CJC
- Gavrus R, see Acalovschi I
- Geary BT, see Nielsen VG
- Gelb AW, see Cechetto DF
- Gerancher JC, see Dolinski SY
- Gerber A, see Mönkhoff M
- Gerber C, see Borgeat A
- Gerber M, see Wahr JA
- Gerig HJ, see Heidegger T
- Ger L-P, see Ho S-T
- Ghaleb AH, see Khorasani A
- Ghobashy AM, see Awad AA
- Ghonaime EA, see White PF
- Gibson CJ, see Cechetto DF
- Gillham MJ, Tousignant CP. Diagnosis by intraoperative transesophageal echocardiography of acute thrombosis of mechanical aortic valve prosthesis associated with the use of biological glue, 92:1123
- Gin T, see Chen PP
- Gintautas J, see Doss NW
- Gippner-Steppert C, see Royston D
- Girling KJ, Spendlove JL, Quraishi MS, Mahajan RP. Video imaging to assess neuromuscular blockade at the larynx, 92:149
- Glass PSA, see Gan TJ
- Godet G
- Against the discontinuation of angiotensin II antagonists before surgery (letter, reply), 92:1617
- see Bertrand M
- Goeters C, see Brodner G
- Golkaryeh M, see Jahr JS
- Goll V, Akça O, Greif R, Freitag H, Arkilic CF, Scheck T, Zoeggeler A, Kurz A, Krieger G, Lenhardt R, Sessler DI. Ondansetron is no more effective than supplemental intraoperative oxygen for prevention of postoperative nausea and vomiting, 92:112
- Gomola A, see Dieudonne N
- Goodchild CS, see Wadhwa A
- Goodman E, Johnson PA. End-tidal carbon

- dioxide tracing configuration depends on sampling size (letter), 92:1357
- Gorton HJ, see Robinson AP
- Goto E, see Tanigawa K
- Goto F, see Usui T
- Gottrup H, see Nikolajsen L
- Gottschalk LI, see Pivalizza EG
- Gottschall JL, see Wohlfeil ER
- Gottumukkala VNR, see Philip J
- Goursot G, see Junca A
- Gouverneur J-MA, see Singelyn FJ
- Goyer C, see Bernard F
- Gravlee GP, see Heres EK
- Gray PE, Perrino AC Jr. Hemodynamic-induced changes in aortic valve area: implications for Doppler cardiac output determinations, 92:584
- Graziano GP, see Samra SK
- Greengrass RA, Klein SM
- Gregory G, see Russell IA
- Greher M, see Marhofer P; Retzl G
- Greif R, see Goll V
- Greilich PE
- Virella CD, Rich JM, Kurada M, Roberts K, Warren JF, Harford WV. Remifentanyl versus meperidine for monitored anesthesia care: a comparison study in older patients undergoing ambulatory colonoscopy, 92:80
- see Sun Y
- Greim C-A, Trautner H, Krämer K, Zimmermann P, Apfel CC, Roewer N. The detection of interatrial flow patency in awake and anesthetized patients: a comparative study using transnasal transesophageal echocardiography, 92:1111
- Gretenkort P, see Harke H
- Grimaud D, see Carles M
- Groban L, Deal DD, Vernon JC, James RL, Butterworth J. Cardiac resuscitation after incremental overdosage with lidocaine, bupivacaine, levobupivacaine, and ropivacaine in anesthetized dogs, 92:37
- Grocott HP, see Newman MF
- Grond S, Demopulos G, Herz J, Palmer PP. Inhibition of synovial plasma extravasation by preemptive administration of an antiinflammatory irrigation solution in the rat knee, 92:1301
- Gross A, see Fischer LG
- Gross ER, Gare M, Toller WG, Kersten JR, Warltier DC, Pagel PS. Ethanol enhances the functional recovery of stunned myocardium independent of K_{ATP} channels in dogs, 92:299
- Gross RM, see Dimaculangan DP
- Gruber EM
- Laussen PC, Casta A, Zimmerman AA, Zurakowski D, Reid R, Odegard KC, Chakravorti S, Davis PJ, McGowan FX Jr, Hickey PR, Hansen DD. Stress response in infants undergoing cardiac surgery: a randomized study of fentanyl bolus, fentanyl infusion, and fentanyl-midazolam infusion, 92:882
- Tschernko EM, Kritzing M, Deviatko E, Wisser W, Zurakowski D, Haider W. The effects of thoracic epidural analgesia with bupivacaine 0.25% on ventilatory mechanics in patients with severe chronic obstructive pulmonary disease, 92:1015
- Gupta S, Singh B, Kachru N. "Blocked" epidural catheter: another cause (letter), 92:1617
- Gustorff B
- Felleiter P, Nahlik G, Brannath W, Hoerauf KH, Spacek A, Kress HG. The effect of remifentanyl on the heat pain threshold in volunteers, 92:369
- see Franz A
- Guyatt GH, see Pronovost PJ
- Gyermek L, see Nishiyama T
- Haas DA, see Wong JK
- Hadani M, see Berkenstadt H
- Hadžić A, see Vloka JD
- Haeseler G, Störmer M, Bufler J, Dengler R, Hecker H, Piepenbrock S, Leuwer M. Propofol blocks human skeletal muscle sodium channels in a voltage-dependent manner, 92:1192
- Hagl C, see Sekaran P
- Hahnenkamp K, see Strunk V
- Haider W, see Gruber EM
- Haisch G, Boldt J, Krebs C, Kumle B, Suttner S, Schulz A. The influence of intravascular volume therapy with a new hydroxyethyl starch preparation (6% HES 130/0.4) on coagulation in patients undergoing major abdominal surgery, 92:565
- Hallén J, see Rawal N
- Halpern SH, see Choi PT-L; Morgan PJ
- Hamaya Y, see Yanagidate F
- Hamer H, see Koessler MJ
- Hammer GB. Pediatric thoracic anesthesia, 92:1449
- Hamza MA, see White PF
- Hanaoka K, see Nishiyama T; Orii R
- Hanazaki M, Jones KA, Perkins WJ, Warner DO. The effects of ethanol on Ca^{2+} sensitivity in airway smooth muscle, 92:767
- Hanley FL, see Russell IA
- Hannallah RS, see Finkel JC; Patel RI
- Hansen DD, see Gruber EM
- Hanusa C, see Hemmerling TM
- Hao S, Takahata O, Iwasaki H. Antinociceptive interaction between spinal clonidine and lidocaine in the rat formalin test: an isobolographic analysis, 92:733
- Hara T, Tomiyasu S, Sungsam C, Fukusaki M, Sumikawa K. Sevoflurane protects stunned myocardium through activation of mitochondrial ATP-sensitive potassium channels, 92:1139
- Hardman JG, see Bedforth N; Bowie RA
- Harford WV, see Greilich PE
- Harke H, Gretenkort P, Ladleif HU, Rahman S, Harke O. The response of neuropathic pain and pain in complex regional pain syndrome I to carbamazepine and sustained-release morphine in patients pretreated with spinal cord stimulation: a double-blinded randomized study, 92:488
- Harke O, see Harke H
- Harks A, see Brodner G
- Harnett MJP, Datta S, Bhavani-Shankar K. The effect of magnesium on coagulation in parturients with preeclampsia, 92:1257
- Harrington D, see Leyvi G
- Harth M, see Wiesner G
- Hartmann B, see Junger A
- Hartmann M, see Schwarzkopf KRG
- Hashimoto S, see Tsuda A
- Hayashida M, see Orii R
- Hayashi Y, see Kariya N
- Haygood RM, see Hord AH
- Hazama K, see Omote K
- Heath BJ, see Mychaskiw G II
- Hebl JR
- Horlocker TT, Pritchard DJ. Diffuse brachial plexopathy after interscalene blockade in a patient receiving cisplatin chemotherapy: the pharmacologic double crush syndrome, 92:249
- Rose SH, Narr BJ, Rorie DK. Postoperative left vocal cord dysfunction caused by Ortner's cardiopharyngeal syndrome, 92:1071
- Hecker H, see Haeseler G
- Hedenstierna G, see Neumann P

- Heidegger T, Gerig HJ, Ulrich B, Kreienbühl G. Validation of a simple algorithm for tracheal intubation: daily practice is the key to success in emergencies—an analysis of 13,248 intubations, 92:517
- Heimann K, see Bremerich DH
- Helfritsch H, see Schwarzkopf K
- Helmy SAK, Bali A. The effect of the preemptive use of the NMDA receptor antagonist dextromethorphan on postoperative analgesic requirements, 92:739
- Hemmerling TM
- Schmidt J, Jacobi KE, Klein P. Intraoperative monitoring of the recurrent laryngeal nerve during single-lung ventilation in esophagectomy, 92:662
- Schmidt J, Wolf T, Hanusa C, Siebzebruehl E, Schmitt H. Intramuscular versus surface electromyography of the diaphragm for determining neuromuscular blockade, 92:106
- Hempelmann G, see Engel JM; Junger A
- Henzi I, see Walder B
- Heres EK, Speight K, Benckart D, Marquez J, Gravlee GP. The clinical onset of heparin is rapid, 92:1391
- Hering R, Wrigge H, Vorwerk R, Brensing KA, Schröder S, Zinserling J, Hoeft A, Spiegel TV, Putensen C. The effects of prone positioning on intraabdominal pressure and cardiovascular and renal function in patients with acute lung injury, 92:1226
- Hernández-Palazón J, Tortosa JA, Martínez-Lage JF, Pérez-Flores D. Intravenous administration of propacetamol reduces morphine consumption after spinal fusion surgery, 92:1473
- Hershman R, see Ben-David B
- Hertle L, see Brodner G
- Herz J, see Grond S
- Heurteaux C, see Lang-Lazdunski L
- Heylen R, see Vranken JH
- Hickey PR, see Gruber EM
- Hickey RF, see Russell IA
- Higashizawa T, see Yuasa H
- Higuchi H
- Adachi Y, Wada H, Kanno M, Satoh T. The effects of low-flow sevoflurane and isoflurane anesthesia on renal function in patients with stable moderate renal insufficiency, 92:650
- Composition of CO₂ absorbents (letter, reply), 92:1357
- see Adachi YU
- Hillel Z, see Cheng DCH; Howie MB
- Himmelseher S
- Pfenninger E, Werner C for the Scientific Neuroanesthesia Research Group of the German Society of Anesthesia and Intensive Care Medicine. Intraoperative monitoring in neuroanesthesia: a national comparison between two surveys in Germany in 1991 and 1997, 92:166
- Ziegler-Pithamitsis D, Argiriadou H, Martin J, Jelen-Esselborn S, Kochs E. Small-dose S(+)-ketamine reduces postoperative pain when applied with ropivacaine in epidural anesthesia for total knee arthroplasty, 92:1290
- Hinder F, see Booke M
- Hirakawa M, see Yokoyama M
- Hirata K, see Fujioka S
- Hirose T, see Asai T
- Höchtel A, see Marhofer P
- Hodgson PS, Liu SS. A comparison of ropivacaine with fentanyl to bupivacaine with fentanyl for postoperative patient-controlled epidural analgesia, 92:1024
- Hoeft A, see Hering R
- Hoenemann CW, see Hollmann MW; Liu H-T
- Hoerauf K, see Wiesner G
- Hoerauf KH, see Gustorff B
- Hoff H, see Schwarzkopf KRG
- Hoffmeyer P, see Culebras X
- Hogue C, see Cheng DCH; Howie MB
- Hohaus T, see Engel JM
- Ho KM, Lim HH. Absence of reflex tachycardia after spinal anesthesia in the elderly (letter, reply), 92:1357
- Hollmann MW
- Liu H-T, Hoenemann CW, Liu W-H, Durieux ME. Modulation of NMDA receptor function by ketamine and magnesium. Part II: interactions with volatile anesthetics, 92:1182
- see Fischer LG; Liu H-T
- Holzman RS, Riley LE, Aron E, Fetherston J. Perioperative care of a patient with acute fatty liver of pregnancy, 92:1268
- Hopkins R, see Maslow A
- Hord AH
- Denson DD, Stowe B, Haygood RM. α -1 and α -2 adrenergic antagonists relieve thermal hyperalgesia in experimental mononeuropathy from chronic constriction injury, 92:1558
- see Kaushik S
- Horlocker TT, see Hebl JR
- Horowitz M, see Mychaskiw G II
- Hoshi T, see Fujii Y
- Hossain S, see Leyvi G
- Ho S-T, Wang J-J, Tzeng J-I, Liu H-S, Ger L-P, Liaw W-J. Dexmethasone for preventing nausea and vomiting associated with epidural morphine: a dose-ranging study, 92:745
- Howie M, see Cheng DCH
- Howie MB
- Cheng D, Newman MF, Pierce ET, Hogue C, Hillel Z, Bowdle TA, Bukunya D. A randomized double-blinded multicenter comparison of remifentanyl versus fentanyl when combined with isoflurane/propofol for early extubation in coronary artery bypass graft surgery, 92:1084
- Dzwonczyk R, McSweeney TD. An evaluation of a new two-electrode myocardial electrical impedance monitor for detecting myocardial ischemia, 92:12
- Huang EY-K, see Li J-Y
- Huber PJ Jr, see Coloma M
- Huizar KM, see Finucane BT
- Hu JW, see Wong JK
- Hummer KA, see Finkel JC
- Hunt JO, see Myles PS
- Huraux C, Ankri A, Eyraud D, Sevin O, Ménégau F, Coriat P, Samama C-M. Hemostatic changes in patients receiving hydroxyethyl starch: the influence of ABO blood group, 92:1396
- Hynninen M, Borger MA, Rao V, Weisel RD, Christakis GT, Carroll J-A, Cheng DCH. The effect of insulin cardioplegia on atrial fibrillation after high-risk coronary bypass surgery: a double-blinded, randomized, controlled trial, 92:810
- Iamaroon A, Pitimana-aree S, Prechawai C, Anusit J, Somcharoen K, Chaiyaroj O. Endotracheal intubation with thiopental/succinylcholine or sevoflurane-nitrous oxide anesthesia in adults: a comparative study, 92:523
- Ide T, Okitsu Y, Nishihashi S, Yamamoto F, Nishino T. The effect of epidural anesthesia on respiratory distress induced by airway occlusion in isoflurane-anesthetized cats, 92:749
- Iijima T, Iwao Y, Sankawa H, Nakamura Z. The epileptogenic property of

- sevoflurane and in patients without epilepsy (letter, reply), 92:1359
- Imanaga K, see Wajima Z
- Imanaka H, Shimaoka M, Matsuura N, Nishimura M, Ohta N, Kiyono H. Ventilator-induced lung injury is associated with neutrophil infiltration, macrophage activation, and TGF- β 1 mRNA upregulation in rat lungs, 92:428
- Inada T
- Taniuchi S, Shingu K, Kobayashi Y, Fujisawa J, Nakao S-i. Propofol depressed neutrophil hydrogen peroxide production more than midazolam, whereas adhesion molecule expression was minimally affected by both anesthetics in rats with abdominal sepsis, 92:437
- see Nagata A
- Inoue S, see Yoshitani K
- Inoue T, see Wajima Z
- Ipe J, see Doss NW
- Irefin SA, see Parker BM
- Ishihara H, see Suzuki A
- Ishizawa Y, see Ma H-C
- Ishizu T, see Tokioka H
- Isner JM, see Lathi KG
- Isowaki S, see Nagata E
- Itano Y, see Yokoyama M
- Iwaki T, see Tokioka H
- Iwanaga S, see Tsuda A
- Iwao Y, see Iijima T
- Iwasaki H, see Hao S
- Jackson MR, see Sun Y
- Jacob H, see Borgeat A
- Jacob LP, see Basto ER
- Jacobi KE, see Hemmerling TM
- Jacobs R, see Sekaran P
- Jadad AR, see Choi PT-L
- Jahr JS, Lurie F, Xi S, Golkaryeh M, Kuznetsova O, Kullar R, Driessen B. A novel approach to measuring circulating blood volume: the use of a hemoglobin-based oxygen carrier in a rabbit model, 92:609
- Jalbout MI, see Aouad MT
- James RL, see Butterworth J; Groban L
- Janelle GM, Urdaneta F, Blas ML, Shryock J, Tang Y-S, Martin TD, Lobato EB. Inhibition of phosphodiesterase Type III before aortic cross-clamping preserves intramyocardial cyclic adenosine monophosphate during cardiopulmonary bypass, 92:1377
- Jäntti V, Yli-Hankala A, Vakkuri A. The epileptogenic property of sevoflurane and in patients without epilepsy (letter), 92:1359
- Jantzen EC, see Fazi L
- Jedeikin R, see Fredman B
- Jelen-Esselborn S, see Himmelseher S
- Jensen TS, see Nikolajsen L
- Jeong SW, see Yoo KY
- Jhaveri R, see Gan TJ
- Jiang H, see Chen J
- Jin C-H, see Liu Y-H
- Jochum M, see Royston D
- Johnson D, see Swenson JD
- Johnson ME, see Fiechtner BK
- Johnson PA, see Goodman E
- Johnson R, see Coates KM
- Joh S, see Satoh K-i
- Jones KA, see Hanazaki M
- Jones RH, see Newman MF
- Joo HS, Kapoor S, Rose DK, Naik VN. The intubating laryngeal mask airway after induction of general anesthesia versus awake fiberoptic intubation in patients with difficult airways, 92:1342
- Joseph NJ, see Nimmagadda U
- Jost A, see Junger A
- Jou I-M, see Tsai Y-C
- Jourdain O, see Bernard J-M
- Junca A, Marret E, Goursot G, Mazoit X, Bonnet F. A comparison of ropivacaine and bupivacaine for cervical plexus block, 92:720
- Junger A, Hartmann B, Benson M, Schindler E, Dietrich G, Jost A, Béye-Basse A, Hempelmann G. The use of an anesthesia information management system for prediction of antiemetic rescue treatment at the postanesthesia care unit, 92:1203
- Jürgensen I, see Engel JM
- Jurim O, see Matot I
- Kachru N, see Gupta S
- Kadar AG, Kramer BA, Barth MC, White PF. Rapacuronium: an alternative to succinylcholine for electroconvulsive therapy, 92:1171
- Kain ZN
- Wang S-M, Mayes LC, Krivutza DM, Teague BA. Sensory stimuli and anxiety in children undergoing surgery: a randomized, controlled trial, 92:897
- see Wang S-M
- Kakihana Y, see Nagata E
- Kaku R, see Yokoyama M
- Kalberer F, see Borgeat A
- Kallmeyer JJ, Collard CD, Fox JA, Body SC, Shernan SK. The safety of intraoperative transesophageal echocardiography: a case series of 7200 cardiac surgical patients, 92:1126
- Kamenik M, Paver-Erzen V. The effects of lactated Ringer's solution infusion on cardiac output changes after spinal anesthesia, 92:710
- Kanaya N
- Nakayama M, Seki S, Kawana S, Watanabe H, Namiki A. Two-person technique for fiberoptic-aided tracheal intubation in a patient with a long and narrow retropharyngeal air space, 92:1611
- Namiki A. The three-dimensional computed tomography imaging and prediction of unanticipated difficult tracheal intubation (letter, reply), 92:282
- Kanmura Y, see Nagata E
- Kanno M, see Higuchi H
- Kapoor S, see Joo HS
- Kapral S, see Retzl G
- Kapuler V, see Eilig I
- Karaca EP, Schwarzenberger J. Practical Echocardiography: The Heart of the Matter. Part 1, Basics (book review), 92:285
- Karasawa F
- Mori T, Kawatani Y, Ohshima T, Satoh T. Deflationary phenomenon of the nitrous oxide-filled endotracheal tube cuff after cessation of nitrous oxide administration, 92:145
- Mori T, Okuda T, Satoh T. Profile soft-seal cuff, a new endotracheal tube, effectively inhibits an increase in the cuff pressure through high compliance rather than low diffusion of nitrous oxide, 92:140
- Kariger R, see Tang J
- Kariya N, Shindoh M, Hayashi Y, Nakasuji M, Nishi S, Nishikawa K, Asada A. A case of fatal paradoxical fat embolism syndrome detected by intraoperative transesophageal echocardiography, 92:688
- Karlan MS, see Tang J
- Karlan SR, see Tang J
- Karzai W, see Schwarzkopf K
- Kattapuram M, see Skubas NJ
- Katz JD. Issues of concern for the aging anesthesiologist, 92:1487
- Kaufman B, see Bekker AY
- Kaushik S, Hord AH, Denson DD, McAllister DV, Smitra S, Allen MG, Prausnitz MR. Lack of pain associated with microfabricated microneedles, 92:502
- Kawaguchi M, see Yoshitani K

- Kawahito S, Kitahata H, Tanaka K, Nozaki J, Oshita S. Pulmonary arterial pressure can be estimated by transesophageal pulsed Doppler echocardiography, 92:1364
- Kawamata M, see Omote K
- Kawamata T, see Omote K
- Kawamura M, see Masaki E
- Kawana S, see Kanaya N
- Kawasaki C, see Kawasaki T
- Kawasaki T, Ogata M, Kawasaki C, Tomihisa T, Okamoto K, Shigematsu A. Surgical stress induces endotoxin hyporesponsiveness and an early decrease of monocyte mCD14 and HLA-DR expression during surgery, 92:1322
- Kawasaki-Yatsugi S, see Nishiyama T
- Kawatani Y, see Karasawa F
- Kaye AD, Vig S, Sabar R. Cerebral hemodynamic response to the introduction of desflurane: a comparison with sevoflurane (letter), 92:282
- Kee S, see Pivalizza EG
- Kehlet H, see Brodner G
- Keller C, see Brimacombe J
- Kennedy DJ
Basics of Anesthesia, 4th Edition (book review), 92:1360
see Sherman SV
- Kerssens C, Klein J, van der Woerd A, Bonke B. Auditory information processing during adequate propofol anesthesia monitored by electroencephalogram bispectral index, 92:1210
- Kersten JR, see Gross ER
- Kessler P, see Bremerich DH
- Khaw KS, see Ngan Kee WD
- Khorasani A, Candido KD, Ghaleb AH, Saatee S, Appavu SK. Canister tip orientation and residual volume have significant impact on the dose of benzocaine delivered by hurricane spray, 92:379
- Kida H, see Satsumae T
- Kiefer R-T
Eliciting paresthesias for peripheral nerve block: a harmful clinical standard? (letter), 92:795
Klaus U, Wiech K, Birbaumer N. The analgesic potency of NMDA-antagonists—a question of mechanism-based use and timing? (letter), 92:795
- Kim HS, see Yoo KY
- Kim MS, see Finkel JC
- Kimura K, see Yokota K
- Kingery WS, Agashe GS, Sawamura S, Davies MF, Clark JD, Maze M. Glucocorticoid inhibition of neuropathic hyperalgesia and spinal fos expression, 92:476
- Kirchner JL, see Newman MF
- Kissin I, Bright CA, Bradley EL Jr. Can inflammatory pain prevent the development of acute tolerance to alfentanil?, 92:1296
- Kitaguchi K, see Yoshitani K
- Kitahata H, see Kawahito S
- Kiyono H, see Imanaka H
- Kiyoshima T, see Matsumoto M
- Klafta JM. Keep it simple (letter), 92:796
- Klaus U, see Kiefer R-T
- Klein E, see Fredman B
- Klein J, see Kerssens C
- Klein P, see Hemmerling TM
- Klein SM, Greengrass RA, Nielsen K, Steele SM. Difficult insertion of interscalene brachial plexus catheter (letter, reply), 92:1074
- Klein U, see Schwarzkopf K
- Kline B, see McNulty SE
- Kobayashi Y, see Inada T
- Kochs E, see Himmelseher S; Vahle-Hinz C
- Kodama S-i, see Nishina K
- Koessler MJ, Fabiani R, Hamer H, Pitto RP. The clinical relevance of embolic events detected by transesophageal echocardiography during cemented total hip arthroplasty: a randomized clinical trial, 92:49
- Koga Y, see Yuasa H
- Koivusalo A-M, see Laisalmi M
- Kosogabe Y, see Tokioka H
- Kox WJ, see Von Dossow V
- Kozek-Langenecker SA, see Franz A
- Kozloski LJ, Aram L
- Kramer BA, see Kadar AG
- Krämer K, see Greim C-A
- Krane EJ, Aram L
- Krebs C, see Haisch G
- Kreienbühl G, see Heidegger T
- Kress HG, see Gustorff B
- Krieger G, see Goll V
- Krismer AC, Lindner KH, Wenzel V, Mayr VD, Voelckel WG, Lurie KG, Strohmenger HU. The effects of endogenous and exogenous vasopressin during experimental cardiopulmonary resuscitation, 92:1499
- Kristensen AGD, see Nikolajsen L
- Kristoffersson AHM, see Finucane BT
- Kritzinger M, see Gruber EM
- Krivutza DM, see Kain ZN
- Krumm B, see Fischer LG
- Kryger KC, Miller RD. *Anesthesia & Analgesia* online: Internet technology and innovation, 92:1363
- Kuhlman G, see Liu N
- Kullar R, see Jahr JS
- Kumle B, see Haisch G
- Kurada M, see Greilich PE
- Kurth CD, see Fazi L
- Kurz A, see Goll V; Taguchi A
- Kuznetsova O, see Jahr JS
- Labbez F, see Legros CB
- Lachat ML, see Zaugg M
- Ladleif HU, see Harke H
- Lahey SJ, see Surgenor SD
- Laisalmi M, Eriksson H, Koivusalo A-M, Pere P, Rosenberg P, Lindgren L. Ketorolac is not nephrotoxic in connection with sevoflurane anesthesia in patients undergoing breast surgery, 92:1058
- Lambert A-S, see Miller JP
- Lam J, see Berkenbosch JW
- Lamm C, see Fitzgerald RD
- Landru JP, see Basto ER
- Langer C, see Swindoll S
- Langeron O, Doelberg M, Ang E-T, Bonnet F, Capdevila X, Coriat P. Voluven, a lower substituted novel hydroxyethyl starch (HES 130/0.4), causes fewer effects on coagulation in major orthopedic surgery than HES 200/0.5, 92:855
- Lang-Lazdunski L, Heurteaux C, Dupont H, Rouelle D, Widmann C, Mantz J. The effects of FK506 on neurologic and histopathologic outcome after transient spinal cord ischemia induced by aortic cross-clamping in rats, 92:1237
- Lang T, see Mahla E
- Lappas DG, see Skubas NJ
- Laskowitz DT, see Newman MF
- Laster MJ, see Eger EI II
- Lathi KG, Vale PR, Losordo DW, Cespedes RM, Symes JF, Esakof DD, Maysky M, Isner JM. Gene therapy with vascular endothelial growth factor for inoperable coronary artery disease: anesthetic management and results, 92:19
- Laurito CE
Regional Anesthesia; Volume 12, Number 2 of *Problems in Anesthesia* (book review), 92:1078
see Pirec V
- Laussen PC, see Gruber EM

- Lavand'homme P
Veyckemans F, Roelants F. Remifentanyl is a valuable alternative to contraindicated neuraxial analgesia in the parturients (letter), 92:1355
see Singelyn FJ
- Leavitt BJ, see Surgenor SD
Leavitt ML, see Sekaran P
Lebowitz P, see Dexter F
Lecarpentier Y, see David J-S
Lee BB, see Ngan Kee WD
Lee C, see Nishiyama T
Lee JU, see Yoo KY
Legros CB, Orliaguet GA, Mayer M-N, Labbez F, Carli PA. Severe anaphylactic reaction to cisatracurium in a child, 92:648
Lehmann A, Boldt J. Remifentanyl in cardiac surgery (letter), 92:557
Lenhardt R, see Goll V
Leonard-White A, see May JA
Lepri A, see Sia S
Le Roux D, see Bernard J-M
Leslie K, see Watson R
Leuwer M, see Haeseler G
Leveno KJ, see Alexander JM
Levitin RM, see Ochroch EA
Levron J-C, see Liu N
Lewis E, Rogachev B, Shaked G, Douvdevani A. The *in vitro* effects of ketamine at large concentrations can be attributed to a nonspecific cytostatic effect, 92:927
Leyvi G, Shore-Lesserson L, Harrington D, Vela-Cantos F, Hossain S. An investigation of a new activated clotting time MAX-ACT in patients undergoing extracorporeal circulation, 92:578
Liaudet L, see Broccard AF
Liaw W-J, see Ho S-T
Li J-Y, Wong C-H, Huang EY-K, Lin Y-C, Chen Y-L, Tan PPC, Chen J-C. Modulations of spinal serotonin activity affect the development of morphine tolerance, 92:1563
Lim HH, see Ho KM
Lim Y-J, see Bahk J-H
Lin C-J, see Yeh H-M
Lin C-R, see Tan P-H
Lindahl SGE, see Modalen ÅÖ
Lindgren L, see Laisalmi M
Lindinger W, see Rieder J
Lindner KH, see Krismer AC; Voelckel WG
Lin YA, see Butterworth J
Lin Y-C, see Li J-Y
Lirk P, see Rieder J
Li S, see White PF
Liu H-S, see Ho S-T
Liu H-T
Hollmann MW, Liu W-H, Hoenemann CW, Durieux ME. Modulation of NMDA receptor function by ketamine and magnesium: Part I, 92:1173
see Hollmann MW
Liu JYW, see Ngan Kee WD
Liu K, see Tan P-H
Liu N, Kuhlman G, Dalibon N, Moutafis M, Levron J-C, Fischler M. A randomized, double-blinded comparison of intrathecal morphine, sufentanil and their combination versus IV morphine patient-controlled analgesia for postthoracotomy pain, 92:31
Liu SS, see Hodgson PS
Liu W-H, see Hollmann MW; Liu H-T
Liu Y-H, Wang J-J, Chang C-F, Jin C-H. Difficult tracheal intubation as a result of unsuspected abnormality of the temporomandibular joint, 92:783
Lobato E, see Blas ML
Lobato EB, see Janelle GM
Lokhandwala BS, see Dimaculangan DP
Loring SH, see Maslow A
Loskota W, see Zárate E
Losordo DW, see Lathi KG
Lotan D, see Vaknin Z
Lovely LK, see Samra SK
Lozada LJ, see Munis JR
Lubarsky DA, see Dexter F; Gan TJ
Lu C-Y, see Tan P-H
Lurie F, see Jahr JS
Lurie KG, see Krismer AC; Voelckel WG
Lu Y, see Pirec V
Lynch JP, see Tung A
Lyons GR, see Robinson AP
Macario A, see Dexter F
Macchi P, see Carles M
MacDonald J, see Swenson JD
MacKinnon S, see Maslow A
MacPherson R. Current Issues in Anesthesiology; Volume 38, Number 4 of *International Anesthesiology Clinics*, 92:1620
Madan R, see Gan TJ
Magistris L, see Casati A
Magotteaux V, see Vergnion M
Mahajan RP, see Bowie RA; Girling KJ
Ma H-C, Dohi S, Wang Y-F, Ishizawa Y, Yanagidate F. The antinociceptive and sedative effects of carbachol and oxycodone administered into brainstem pontine reticular formation and spinal subarachnoid space in rats, 92:1307
Mahla E, Lang T, Vicenzi MN, Werkgartner G, Maier R, Probst C, Metzler H. Thromboelastography for monitoring prolonged hypercoagulability after major abdominal surgery, 92:572
Maier R, see Mahla E
Makita T, see Tsuda A
Makuuchi M, see Orii R
Malik N, see Choi PT-L
Ma M, see Chen PP
Mandell G, see Fischer LG
Mandell GL, see Vallejo MC
Manisterski Y, see Vaknin Z
Manninen PH, see Blanshard HJ
Mantz J, see Lang-Lazdunski L
Marchini O, see Sia S
Margalit N, see Berkenstadt H
Margarit S, see Acalovschi I
Marhofer P, Freitag H, Höchtel A, Greher M, Erlacher W, Semsroth M. S(+)-ketamine for rectal premedication in children, 92:62
Marquez J, see Heres EK
Marret E, see Junca A
Marsh HM, see Dexter F
Martens-Lobenhoffer J, Eisenhardt S, Tröger U, Röse W, Meyer FP. The effect of anxiety and personality on the pharmacokinetics of oral midazolam, 92:621
Martin E, see Von Dossow V
Martinez EA, see Pronovost PJ
Martinez-Lage JF, see Hernández-Palazón J
Martin J, see Himmelseher S
Martin TD, see Janelle GM
Masaki E, Yamazaki K, Ohno Y, Nishi H, Matsumoto Y, Kawamura M. The anesthetic interaction between adenosine triphosphate and N-methyl-D-aspartate receptor antagonists in the rat, 92:134
Maslow A, Stearns G, Bert A, Feng W, Price D, Schwartz C, MacKinnon S, Rotenberg F, Hopkins R, Cooper G, Singh A, Loring SH. Monitoring end-tidal carbon dioxide during weaning from cardiopulmonary bypass in patients without significant lung disease, 92:306
Masuzawa M, see Nagata A
Mather LE, see Coates KM
Matot I, Jurim O. The protective effect of acadesine on lung ischemia-reperfusion injury, 92:590
Matsuki A, see Suzuki A
Matsumoto M, Ohtake K, Wakamatsu H, Oka S, Kiyoshima T, Nakakimura

- K, Sakabe T. The time course of acquisition of ischemic tolerance and induction of heat shock protein 70 after a brief period of ischemia in the spinal cord in rabbits, 92:418
- Matsumoto Y, see Masaki E
- Matsuura N, see Imanaka H
- Mauritz W, see Retzl G
- Mayer M-N, see Legros CB
- Mayes LC, see Kain ZN
- May JA, White HC, Leonard-White A, Warltier DC, Pagel PS. The patient recovering from alcohol or drug addiction: special issues for the anesthesiologist, 92:1601
- Mayo A, see Fredman B
- Mayr VD, see Krismer AC
- Maysky M, see Lathi KG
- Maze M, see Kingery WS
- Mazoit X, see Junca A
- Mazze RI. Composition of CO₂ absorbents (letter), 92:1356
- McAllister DV, see Kaushik S
- McCullough JN, see Sekaran P
- McGowan FX Jr, see Gruber EM
- McIntire DD, see Alexander JM
- McKenzie ED, see Stayer SA
- McKnite S, see Voelckel WG
- McNulty SE, Kline B, Welsh J, Bartkowski R. Radiofrequency transmission to monitoring devices in the operating room: a simulation study, 92:384
- McSweeney TD, see Howie MB
- Meersschaert K, see Bertrand M
- Mendelson WB, see Tung A
- Ménégaux F, see Huraux C
- Menges T, see Engel JM
- Mentzelopoulos SD, Tzoufi MJ, Papageorgiou EP. The disposition of the cervical spine and deformation of available cord space with conventional- and balloon laryngoscopy-guided laryngeal intubation: a comparative study, 92:1331
- Merritt WT, see Pronovost PJ
- Mesa A, see Soto R
- Metais C, see Park KW
- Metzler H, see Mahla E
- Metz RJ, see Mortero RF
- Meyer BJ, see Bannister CF
- Meyer FP, see Martens-Lobenhoffer J
- Michael R, see Doss NW
- Michel C, see Bernard J-M
- Michel TC, Rosenberg AL, Polley LS. Obstetric anesthetic management of a parturient with Larsen syndrome and short stature, 92:1266
- Mikawa K, see Nishina K; Shiga M
- Mikhail M, see Roffey P
- Miller Hance WC, see Russell IA
- Miller JP, Lambert A-S, Shapiro WA, Russell IA, Schiller NB, Cahalan MK. The adequacy of basic intraoperative transesophageal echocardiography performed by experienced anesthesiologists, 92:1103
- Miller RD, see Kryger KC
- Mingus M, see Zárate E
- Mirakhur RK, see Elliott P
- Miyabe M, see Fukuda T
- Miyamoto E, see Nagata A
- Mizobuchi S, see Yokoyama M
- Mizutani T, see Fukuda T
- Modalen ÅÖ, Arlander E, Eriksson LI, Lindahl SGE. The effects on hypercarbic ventilatory response of sameridine compared to morphine and placebo, 92:529
- Mohan Reddy V, see Russell IA
- Mok MS, see Yeh H-M
- Mönkhoff M, Schwarz U, Gerber A, Fanconi S, Bänziger O. The effects of sevoflurane and halothane anesthesia on cerebral blood flow velocity in children, 92:891
- Montes FR. Immediate or early extubation: where do we start? (letter, reply), 92:1074
- Morgan PJ, Halpern SH, Tarshis J. The effects of an increase of central blood volume before spinal anesthesia for cesarean delivery: a qualitative systematic review, 92:997
- Mori T, see Karasawa F
- Mortero RF
- Clark LD, Tolan MM, Metz RJ, Tsueda K, Sheppard RA. The effects of small-dose ketamine on propofol sedation: respiration, postoperative mood, perception, cognition, and pain, 92:1465
- Orahovac Z, Tsueda K, Bumpous JM. Severe laryngospasm at tracheal extubation in a patient with superior laryngeal nerve injury, 92:271
- Mourey FD, see Basto ER
- Moutafis M, see Liu N
- Moysa GL, see O'Connor PJ
- Mulroy MF. Ambulatory anesthesia for knee arthroscopy (letter, reply), 92:556
- Munis JR, Bhatia S, Lozada LJ. Peripheral venous pressure as a hemodynamic variable in neurosurgical patients, 92:172
- Muñoz L, see Fernández-Guisasaola J
- Murao K, see Nagata A
- Murthy HS Rao GSU. Cardiovascular responses to scalp infiltration with different concentrations of epinephrine with or without lidocaine during craniotomy, 92:1516
- Muñoz HR, Dagnino JA. Power analysis with more than two groups (letter), 92:797
- Mychaskiw G II
- Horowitz M, Sachdev V, Heath BJ. Explicit intraoperative recall at a bispectral index of 47, 92:808
- see Blakley O
- Myles PS, Hunt JO, Fletcher H. Measuring health status (quality of recovery?) after anesthesia and surgery (letter), 92:281
- Nagano O, see Yokoyama M
- Nagase K, Ando-Nagase K. Endotoxin augments cerebral hyperemic response to halothane by inducing nitric oxide synthase (letter), 92:1075
- Nagata A, Nakao S-i, Nishizawa N, Masuzawa M, Inada T, Murao K, Miyamoto E, Shingu K. Xenon inhibits but N₂O enhances ketamine-induced c-Fos expression in the rat posterior cingulate and retrosplenial cortices, 92:362
- Nagata E, Kakihana Y, Tobo K, Isowaki S, Kanmura Y. The effects of olprinone (a phosphodiesterase III inhibitor) on hepatic vascular bed in a porcine model of endotoxemia, 92:676
- Naguib M
- The three-dimensional computed tomography imaging and prediction of unanticipated difficult tracheal intubation (letter), 92:281
- see Dexter F
- Nahlik G, see Gustorff B
- Naik VN, see Joo HS
- Nakae Y, Fujita S, Namiki A. Morphine enhances myofilament Ca²⁺ sensitivity in intact guinea pig beating hearts, 92:602
- Nakajo N, see Tomioka S
- Nakakimura K, see Matsumoto M
- Nakamura Y, see Tokioka H
- Nakamura Z, see Iijima T
- Nakao S-i, see Inada T; Nagata A
- Nakasuiji M, see Kariya N
- Nakatsuka H, see Yokoyama M
- Nakayama H, see Fukuda T
- Nakayama M, see Kanaya N
- Nakayama Y, see Omote K

- Namiki A, see Kanaya N; Nakae Y; Omote K
- Narr BJ, see Hebl JR
- Naruse R, see Tang J
- Nash HA, see Walcourt A
- Nathanson M, see Bedford N
- Nehashi S, see Ide T
- Neidhardt B, see Schmitt HJ
- Neidhardt G, see Bremerich DH
- Nelson TE, see Zapata-Sudo G
- Neumann P, Hedenstierna G. Ventilatory support by continuous positive airway pressure breathing improves gas exchange as compared with partial ventilatory support with airway pressure release ventilation, 92:950
- Neuvonen PT, van den Berg AA. Postoperative coma in a child with carnitine palmitoyltransferase I deficiency, 92:646
- Nevin K, see Finucane BT
- Newman MF
- Laskowitz DT, White WD, Kirchner JL, Grocott HP, Stafford-Smith M, Sketch MH, Jones RH, Reves JG, Saunders AM. Apolipoprotein E polymorphisms and age at first coronary artery bypass graft, 92:824
- see Cheng DCH; Howie MB
- Ngan Kee WD, Khaw KS, Lee BB, Wong ELY, Liu JYW. The limitations of ropivacaine with epinephrine as an epidural test dose in parturients, 92:1529
- Nielsen K, Klein SM
- Nielsen VG
- Armstead VE, Geary BT, Opentanova IL. Pentalyte does not decrease heparinoid release but does decrease circulating thrombotic mediator activity associated with aortic occlusion-reperfusion in rabbits, 92:314
- Nitric oxide decreases coagulation protein function in rabbits as assessed by thromboelastography, 92:320
- see Chaney JD
- Nikolajsen L, Gotttrup H, Kristensen AGD, Jensen TS. The analgesic potency of NMDA-antagonists—a question of mechanism-based use and timing? (letter, reply), 92:795
- Nimmagadda U, Chiravuri SD, Salem MR, Joseph NJ, Wafai Y, Crystal GJ, El-Orbany MI. Preoxygenation with tidal volume and deep breathing techniques: the impact of duration of breathing and fresh gas flow, 92:1337
- Nishi H, see Masaki E
- Nishikawa K, see Kariya N
- Nishikawa T, see Tanaka M
- Nishimura M, see Imanaka H
- Nishina K
- Mikawa K, Kodama S-i, Obara H. ONO1714, a new inducible nitric oxide synthase inhibitor, attenuates sepsis-induced diaphragmatic dysfunction in hamsters, 92:959
- see Shiga M
- Nishino T, see Ide T
- Nishi S, see Kariya N
- Nishiyama T, Gyermek L, Lee C, Kawasaki-Yatsugi S, Yamaguchi T, Hanaoka K. The analgesic interaction between intrathecal clonidine and glutamate receptor antagonists on thermal and formalin-induced pain in rats, 92:725
- Nishizawa N, see Nagata A
- Nitta R, see Tanaka M
- Nivoche Y, see Cucchiari G
- Novick AC, see O'Hara JF Jr
- Nozaki J, see Kawahito S
- Nugent WC, see Surgenor SD
- Nuttall GA, see Fiechtner BK
- Obara H, see Nishina K; Shiga M
- Ochroch A. Airway Management; Volume 38, Number 3 (Summer 2000) of *International Anesthesiology Clinics* (book review), 92:798
- Ochroch EA
- Barnett RA, Tobias MD. A complication of venous double cannulation (letter), 92:558
- Levitan RM. A videographic analysis of laryngeal exposure comparing the articulating laryngoscope and external laryngeal manipulation, 92:267
- O'Connor CJ, Tuman KJ. Antifibrinolytic therapy and pulmonary thromboembolism during orthotopic liver transplantation (letter, reply), 92:560
- O'Connor GT, see Surgenor SD
- O'Connor PJ
- Moysa GL, Finucane BT. Thoracic epidural anesthesia for bilateral reduction mammoplasty in a patient with Klippel-Feil syndrome, 92:514
- see Bowie RA
- Oczenski W, see Fitzgerald RD
- Odegard KC, see Gruber EM
- Ogata M, see Kawasaki T
- Ogawa R, see Wajima Z
- Ogura A, see Wajima Z
- O'Hara JF Jr, Colburn WA, Tetzlaff JE, Novick AC, Angermeier KW, Schubert A. Hemoglobin and methemoglobin concentrations after large-dose infusions of dapsirin cross-linked hemoglobin, 92:44
- O'Hara P, see Sprung J
- Ohlsson T, see Rawal N
- Ohno Y, see Masaki E
- Ohshima N, see Fukuda T
- Ohshima T, see Karasawa F
- Ohshita N, see Tomioka S
- Ohtake K, see Matsumoto M
- Ohta N, see Imanaka H
- Okada K, see Fujioka S
- Okamoto K, see Kawasaki T
- Oka S, see Matsumoto M
- Okawa H, see Suzuki A
- Okitsu Y, see Ide T
- Okuda T, see Karasawa F
- Oliver WC Jr, see Fiechtner BK
- Olufolabi AJ, see Reynolds JD
- Omote K, Kawamata T, Nakayama Y, Kawamata M, Hazama K, Namiki A. The effects of peripheral administration of a novel selective antagonist for prostaglandin E receptor subtype EP₁, ONO-8711, in a rat model of postoperative pain, 92:233
- Ong BY, see Finucane BT
- Opentanova IL, see Nielsen VG
- Orahovac Z, see Mortero RF
- Orii R, Sugawara Y, Hayashida M, Uchida K, Yamada Y, Takayama T, Makuuchi M, Hanaoka K. Lactate is correlated with the indocyanine green elimination rate in liver resection for cirrhotic patients, 92:1064
- Orliaguet GA, see Legros CB
- Oshita S, see Kawahito S
- Ouda W, see Awad AA
- Oyen LJ, see Fiechtner BK
- Ozier YM, see Dieudonne N
- Paech MJ, see Pavy TJG
- Pagel PS, see Gross ER; May JA
- Paiste J, Williams JP. Unsuccessful placement of transesophageal echocardiography probe because of esophageal pathology, 92:870
- Palmer CM. Continuous intrathecal sufentanil for postoperative analgesia, 92:244
- Palmer PP, see Grond S
- Papageorgiou EP, see Mentzelopoulos SD
- Paret G, see Vaknin Z

- Parker BM, Irefin SA. Antifibrinolytic therapy and pulmonary thromboembolism during orthotopic liver transplantation (letter), 92:559
- Park KW, Metais C, Dai HB, Comunale ME, Sellke FW. Microvascular endothelial dysfunction and its mechanism in a rat model of subarachnoid hemorrhage, 92:990
- Pasque M, see Skubas NJ
- Patel KM, see Finkel JC; Patel RI
- Patel RI, Verghese ST, Hannallah RS, Aregawi A, Patel KM. Fast-tracking children after ambulatory surgery, 92:918
- Paut O, Sallabery M, Schreiber-Deturmeny E, Rémond C, Bruguerolle B, Camboulives J. Continuous fascia iliaca compartment block in children: a prospective evaluation of plasma bupivacaine concentrations, pain scores, and side effects, 92:1159
- Paver-Erzen V, see Kamenik M
- Pavy TJG, Paech MJ, Evans SF. The effect of intravenous ketorolac on opioid requirement and pain after cesarean delivery, 92:1010
- Peeters-Asdourian C, see Simopoulos T
- Pena M, see Finkel JC
- Peng C-H, see Tan P-H
- Penning DH, see Dexter F
- Perel A, see Berkenstadt H
- Pere P, see Laisalmi M
- Perez BJ, see Philip J
- Pérez-Flores D, see Hernández-Palazón J
- Perkins WJ, see Hanazaki M
- Pernia A, see Calderón E
- Perrino AC Jr., see Gray PE
- Peterfreund RA, see Fitzsimons MG
- Pfammatter T, see Zaugg M
- Pfenninger E, see Himmelseher S
- Philip J, Sharma SK, Gottumukkala VNR, Perez BJ, Slaymaker EA, Wiley J. Transient neurologic symptoms after spinal anesthesia with lidocaine in obstetric patients, 92:405
- Piepenbrock S, see Haeseler G
- Pierce E, see Cheng DCH
- Pierce ET, see Howie MB
- Pirec V, Laurito CE, Lu Y, Yeomans DC. The combined effects of N-type calcium channel blockers and morphine on Aδ versus C fiber mediated nociception, 92:239
- Pitimana-aree S, see Iamaroon A
- Pitto RP, see Koessler MJ
- Pivalizza EG, Pivalizza PJ, Kee S, Gottschalk LI, Szmuk P, Abramson DC. Sonoclot analysis in healthy children, 92:904
- Pivalizza PJ, see Pivalizza EG
- Plaisance P, see Voelckel WG
- Plante P, see Atallah F
- Plaza A, see Fernández-Guisasola J
- Pollard JB. Cardiac arrest during spinal anesthesia: common mechanisms and strategies for prevention, 92:252
- Polley LS, see Columb MO; Michel TC
- Ponczecchi P, see Sia S
- Poole W, see Wohlfeil ER
- Popp H, see Sprung J
- Porth J, see Swenson JD
- Prausnitz MR, see Kaushik S
- Prazeller P, see Rieder J
- Prechawai C, see Iamaroon A
- Préfaut C, see Ayoub J
- Pregel AW, Rembecki M, Wenzel V, Steinbach G. A comparison of the endotracheal tube and the laryngeal mask airway as a route for endobronchial lidocaine administration, 92:1505
- Preußler N-P, see Schwarzkopf K
- Price D, see Maslow A
- Pridham JN, see Finucane BT
- Prielipp RC, see Butterworth J
- Prioux J, see Ayoub J
- Pritchard DJ, see Hebl JR
- Probst C, see Mahla E
- Pronovost PJ, Berenholtz SM, Dorman T, Merritt WT, Martinez EA, Guyatt GH. Evidence-based medicine in anesthesiology, 92:787
- Pulcini A, see Carles M
- Putensen C, see Hering R
- Quinn R, see Surgenor SD
- Quraishi MS, see Girling KJ
- Rachinsky M, see Eilig I
- Raeder JC, Steine S, Vatsgar TT. Oral ibuprofen versus paracetamol plus codeine for analgesia after ambulatory surgery, 92:1470
- Rahman S, see Harke H
- Raines DE, see Fitzsimons MG
- Raja SN, see Shir Y
- Rajpal S, see Doss NW
- Ramanathan S, see Vallejo MC
- Ramonatxo M, see Ayoub J
- Rao GSU, see Murthy HS
- Rao V, see Hynninen M
- Raucoules-Aime M, see Carles M
- Rawal N, Allvin R, Amilon A, Ohlsson T, Hallén J. Postoperative analgesia at home after ambulatory hand surgery: a controlled comparison of tramadol, metamizol, and paracetamol, 92:347
- Rehfeldt KH, Cook DJ. Transesophageal echocardiographic imaging of a new aortic cannula for differential perfusion during cardiopulmonary bypass, 92:338
- Reichman C, see Tang J
- Reid R, see Gruber EM
- Rembecki M, see Pregel AW
- Rémond C, Paut O
- Retzl G, Kapral S, Greher M, Mauritz W. Ultrasonographic findings of the axillary part of the brachial plexus, 92:1271
- Reuben SS
- Ambulatory anesthesia for knee arthroscopy (letter), 92:556
- Sklar J, El-Mansouri M. The preemptive analgesic effect of intraarticular bupivacaine and morphine after ambulatory arthroscopic knee surgery, 92:923
- Reves JG, see Newman MF
- Reynolds JD, Booth JV, Olufolabi AJ. A preliminary investigation of remifentanyl as a labor analgesic (letter, reply), 92:1358
- Reynolds LM, see Russell IA
- Rich GF, see Strunk V
- Rich JM, see Greilich PE
- Richter JA, see Dietrich W
- Ricksten S-E, see Thorén A
- Rieder J, Prazeller P, Boehler M, Lirk P, Lindinger W, Amann A. Online monitoring of air quality at the postanesthetic care unit by proton-transfer-reaction mass spectrometry, 92:389
- Riley LE, see Holzman RS
- Riou B, see David J-S
- Rizk LB, see Aouad MT
- Roberts K, see Greilich PE
- Robinette C, see Seitz PA
- Robins B, Boggs DP. Caudal epidural blood patch for treating intractable vomiting in a child after placement of a permanent intrathecal catheter, 92:1169
- Robinson AP, Lyons GR, Wilson RC, Gorton HJ, Columb MO. Levobupivacaine for epidural analgesia in labor: the sparing effect of epidural fentanyl, 92:410
- Roelants F, see Lavand'homme P; Singelyn FJ
- Roewer N, see Greim C-A
- Roffey P, Thangathurai D, Mikhail M. Anesthetic concerns in retroperitoneal lymph node

- dissection for testicular cancer (letter), 92:1618
- Rogachev B, see Lewis E
- Romand JR, see Walder B
- Rorie DK, see Hebl JR
- Rose DK, see Joo HS
- Rose JB, see Fazi L
- Rosenberg AL, see Michel TC
- Rosenberg P, see Laisalmi M
- Rosenthal JA. Atlas of Pain Management Injection Techniques (book review), 92:284
- Rose SH, see Hebl JR
- Röse W, see Martens-Lobenhoffer J
- Ross N, see Davis L
- Rotenberg F, see Maslow A
- Rouelle D, see Lang-Lazdunski L
- Rouine-Rapp K, see Russell IA
- Rowlingson JC. How can local anesthetic in the wound not help?, 92:3
- Royse A, see Royse C
- Royse C, Soeding P, Royse A. Immediate or early extubation: where do we start? (letter), 92:1073
- Royston D
- Cardigan R, Gippner-Steppert C, Jochum M. Is perioperative plasma aprotinin concentration more predictable and constant after a weight-related dose regimen?, 92:830
- see Forrest MR
- Rückert J, see Von Dossow V
- Ruediger HW, see Wiesner G
- Ruetsch YA, see Borgeat A
- Russell IA
- Miller Hance WC, Gregory G, Balea MC, Cassorla L, DeSilva A, Hickey RF, Reynolds LM, Rouine-Rapp K, Hanley FL, Mohan Reddy V, Cahalan MK. The safety and efficacy of sevoflurane anesthesia in infants and children with congenital heart disease, 92:1152
- see Miller JP
- Ruwoldt R, see Engel JM
- Saatee S, see Khorasani A
- Sabar R, see Kaye AD
- Sachdev V, see Mychaskiw G II
- Sacher P, see Simma B
- Sadovnikoff N. Anesthesia and Renal Considerations; Volume 18, Number 4 (December 2000) of *Anesthesiology Clinics of North America*, 92:1360
- Saito M, see Tsuda A
- Saito S, see Usui T
- Saito T. Characteristics of extended unilateral (paravertebral) anesthesia (letter), 92:558
- Sakabe T, see Matsumoto M
- Sakaguchi M, see Satsumae T
- Sakahashi H, see Fujioka S
- Sakamoto N, see Satoh K-i
- Sakamoto T, see Yoshitani K
- Salcedo E, see Bertrand M
- Salem MR, see Nimmagadda U
- Sallabery M, Paut O
- Samama CM
- Thromboelastography: the next step (editorial), 92:563
- see Huraux C
- Samii K, see Atallah F
- Samir H, see Bekker AY
- Samra SK, Dy EA, Welch KB, Lovely LK, Graziano GP. Remifentanyl- and fentanyl-based anesthesia for intraoperative monitoring of somatosensory evoked potentials, 92:1510
- Sankawa H, see Iijima T
- Sari A, see Yokota K
- Sarpal RS, see Fiechtner BK
- Sasaki T, see Fujioka S
- Satoh K-i, Sakamoto N, Shinohe Y, Satoh M, Joh S. Indigo carmine-induced bradycardia in a patient during general anesthesia, 92:276
- Satoh M, see Satoh K-i
- Satoh T, see Adachi YU; Higuchi H; Karasawa F
- Satsumae T, Yamaguchi H, Sakaguchi M, Yasunaga T, Yamashita S, Yamamoto S, Kida H. Preoperative small-dose ketamine prevented tourniquet-induced arterial pressure increase in orthopedic patients under general anesthesia, 92:1286
- Sauer F, see Schwarzkopf K
- Saunders AM, see Newman MF
- Sawamura S, see Kingery WS
- Schaller M-D, see Broccard AF
- Scheck T, see Goll V
- Schiller NB, see Miller JP
- Schindler E, see Junger A
- Schmid ER, see Zaugg M
- Schmidt J, see Hemmerling TM
- Schmitt H, see Hemmerling TM
- Schmitt HJ, Becke K, Neidhardt B. Epidural anesthesia for cesarean section in a patient with polycythemia rubra vera and preeclampsia, 92:1535
- Schneider MC, Birnbach DJ. Lidocaine neurotoxicity in the obstetric patient: is the water safe? (editorial), 92:287
- Schneuing M, see Strunk V
- Schnyder P, see Broccard AF
- Schoenmaekers B, see De Witte JL
- Schreiber-Deturmeny E, see Paut O
- Schreiber SB, see Finkel JC
- Schreiber T, see Schwarzkopf K
- Schröder S, see Hering R
- Schroegendorfer K, see Wiesner G
- Schubert A, see O'Hara JF Jr
- Schulz A, see Haisch G
- Schwartz AJ. Hypertext Book of Pulmonary Artery Catheterization (book review), 92:798
- Schwartz C, see Maslow A
- Schwarzenberger J, see Karaca EP
- Schwarzkopf K, Klein U, Schreiber T, Prießler N-P, Bloos F, Helfritsch H, Sauer F, Karzai W. Oxygenation during one-lung ventilation: the effects of inhaled nitric oxide and increasing levels of inspired fraction of oxygen, 92:842
- Schwarzkopf KRG, Hoff H, Hartmann M, Fritz HG. A comparison between meperidine, clonidine and urapidil in the treatment of postanesthetic shivering, 92:257
- Schwarz U, see Mönkhoff M
- Scott RL, see Walcourt A
- Scuderi P, see Zárate E
- Sebel PS, see Bannister CF
- Seeburger MD, see Zwetsch G
- Segal E, see Berkenstadt H
- Seitz PA, Alfery DD, Robinette C. Acute transverse myelopathy after posterior cervical decompressive laminectomy and fusion, 92:1254
- Sekaran P, Ehrlich M, Hagl C, Leavitt ML, Jacobs R, McCullough JN, Bennett-Guerrero E. A comparison of complete blood replacement with varying hematocrit levels on neurological recovery in a porcine model of profound hypothermic (<5°C) circulatory arrest, 92:329
- Seki S, see Kanaya N
- Sellke FW, see Park KW
- Seltzer Z, see Shir Y
- Semsroth M, see Marhofer P
- Serrano ML, see Fernández-Guisasaola J
- Sessler DI, see De Witte JL; Goll V; Taguchi A
- Sevin O, see Huraux C
- Shaked G, see Lewis E
- Shapira Y, see Eilig I
- Shapiro A, see Fredman B
- Shapiro WA, see Miller JP
- Sharma SK, see Alexander JM; Philip J
- Shelley KH, see Awad AA
- Shepherd AP, see Ali AA
- Sheppard RA, see Mortero RF

- Sherman SV, Wall MH, Kennedy DJ, Brooker RF, Butterworth J. Do pulmonary artery catheters cause or increase tricuspid or pulmonic valvular regurgitation?, 92:1117
- Shernan SK, see Kallmeyer JJ
- Sheth R, see Shir Y
- Shibata O, see Tsuda A
- Shiga M, Nishina K, Mikawa K, Obara H. The effects of lidocaine on nitric oxide production from an activated murine macrophage cell line, 92:128
- Shiga T, see Wajima Z
- Shigematsu A, see Kawasaki T
- Shimaoka M, see Imanaka H
- Shindoh M, see Kariya N
- Shingu K, see Asai T; Inada T; Nagata A
- Shinohe Y, see Satoh K-i
- Shir Y, Sheth R, Campbell JN, Raja SN, Seltzer Z. Soy-containing diet suppresses chronic neuropathic sensory disorders in rats, 92:1029
- Shore-Lesserson L, see Leyvi G
- Shryock J, see Janelle GM
- Shukla RC, see Finucane BT
- Shyu M-K, see Yeh H-M
- Sia S, Bartoli M, Lepri A, Marchini O, Ponzeccchi P. Eliciting paresthesias for peripheral nerve block: a harmful clinical standard? (letter, reply), 92:795
- Siddik SS, see Aouad MT
- Siebzehnruel E, see Hemmerling TM
- Siemers M, see Vahle-Hinz C
- Sigl JC, see Bannister CF
- Silverman DG, see Awad AA
- Simma B, Burger R, Falk M, Sacher P, Torresani T, Fanconi S. The release of antidiuretic hormone is appropriate in response to hypovolemia and/or sodium administration in children with severe head injury: a trial of lactated Ringer's solution versus hypertonic saline, 92:641
- Simopoulos T, Peeters-Asdourian C. Pneumocephalus after cervical epidural steroid injection, 92:1576
- Singelyn FJ. Difficult insertion of interscalene brachial plexus catheter (letter), 92:1074
- Lavand'homme P, De Kock M, Roelants F, Veyckemans F. Regional Anesthesia; Volume 18, Number 2 of *Anesthesiology Clinics of North America* (book review), 92:284
- Vanderelst PE, Gouverneur J-MA. Extended femoral nerve sheath block after total hip arthroplasty: continuous versus patient-controlled techniques, 92:455
- Singh A, see Maslow A
- Singh B, see Gupta S
- Skarvan K, see Zwetsch G
- Sketch MH, see Newman MF
- Sklar J, see Reuben SS
- Skubas NJ, Beardslee M, Barzilai B, Pasque M, Kattapuram M, Lappas DG. Constrictive pericarditis: intraoperative hemodynamic and echocardiographic evaluation of cardiac filling dynamics, 92:1424
- Slaymaker EA, see Philip J
- Sloan F, see Gan TJ
- Sloninsky A, see Tang J
- Smith TC. Proceedings of the Fourth International Symposium on the History of Anaesthesia (book review), 92:1078
- Smitra S, see Kaushik S
- Sobczynski P, see Wiesner G
- Soeding P, see Royse C
- Somcharoen K, see Iamaroon A
- Sonner J, see Zhang Y
- Sonner JM, see Zhang Y
- Soto R, Mesa A. New intravenous catheter not suitable for trans-tracheal jet ventilation (letter), 92:1074
- Spacek A, see Gustorff B
- Spannagl M, see Dietrich W
- Speight K, see Heres EK
- Spendlove JL, see Girling KJ
- Spiegel TV, see Hering R
- Spies CD, see Von Dossow V
- Sprung J, Popp H, O'Hara P, Woletz J. The successful use of hemoglobin-based oxygen carrier as a primary blood substitute during abdominal aneurysm repair with large blood loss, 92:1413
- Stabernack C, see Zhang Y
- Stacey RGW, see Dhileepan S
- Stafford-Smith M, see Newman MF
- Stayer SA. Andropoulos DB, Bent ST, McKenzie ED, Fraser CD. Volume ventilation of infants with congenital heart disease: a comparison of Dräger, NAD 6000 and Siemens, Servo 900C ventilators, 92:76
- Bent ST, Andropoulos DA. Proper probe positioning for infants with compromised ventilation from transesophageal echocardiography (letter), 92:1076
- Stearns G, see Maslow A
- Steele SM, Klein SM
- Steinbach G, see Pregel AW
- Steine S, see Ræder JC
- Steinke JM, see Ali AA
- Stimpfl T, see Fitzgerald RD
- Stix MS. Knees-to-abdomen "Heimlich maneuver" in a morbidly obese patient (letter), 92:1619
- Stonebraker VC, see Ben-David B
- Störmer M, see Haeseler G
- Storm M, see Booke M
- Stout RG, see Awad AA
- Stowe B, see Hord AH
- Strohenger HU, see Krismer AC
- Strunk V, Hahnenkamp K, Schneuing M, Fischer LG, Rich GF. Selective iNOS inhibition prevents hypotension in septic rats while preserving endothelium-dependent vasodilation, 92:681
- Sudo RT, see Zapata-Sudo G
- Sugawara Y, see Orii R
- Sugiyama M, see Yoshitani K
- Sugiyama N, see Yoshitani K
- Sujirattanawimol N, see Fiechtner BK
- Sumikawa K, see Hara T; Tsuda A
- Sungsam C, see Hara T
- Sun W-Z, see Yeh H-M
- Sun Y, Greilich PE, Wilson SIO, Jackson MR, Whitten CW. The use of lepirudin for anticoagulation in patients with heparin-induced thrombocytopenia during major vascular surgery, 92:344
- Surgenor SD, O'Connor GT, Lahey SJ, Quinn R, Charlesworth DC, Dacey LJ, Clough RA, Leavitt BJ, Defoe GR, Fillinger M, Nugent WC for the Northern New England Cardiovascular Disease Study Group. Predicting the risk of death from heart failure after coronary artery bypass graft surgery, 92:596
- Suttner S, see Haisch G
- Suzuki A, Ishihara H, Okawa H, Tsubo T, Matsuki A. Can initial distribution volume of glucose predict hypovolemic hypotension after radical surgery for esophageal cancer?, 92:1146
- Swenson JD, Durcan S, Johnson D, Porth J, MacDonald J. The successful treatment of a spino-subcutaneous fistula after bone marrow harvest by using an epidural blood patch, 92:1050
- Swindoll S, Langer C. Creation of individual soaking compartments within a glutaraldehyde fume hood (letter), 92:1073

- Symes JF, see Lathi KG
 Szmuk P, see Pivalizza EG
 Szpisjak DF. Debris elimination from partially-filled cell salvage bowls, 92:1137
- Taguchi A, Arkilic CF, Ahluwalia A, Sessler DI, Kurz A. Negative pressure rewarming vs. forced air warming in hypothermic postanesthetic volunteers, 92:261
- Taguchi H, see Asai T
 Takahata O, see Hao S
 Takashima T, see Yokoyama M
 Takayama T, see Orii R
 Takazawa A, see Fujioka S
 Takechi M, see Tomioka S
 Takeda T, see Tanigawa K
 Tanaka K, see Kawahito S; Tanigawa K
 Tanaka M
 Nishikawa T. Absence of reflex tachycardia after spinal anesthesia in the elderly (letter), 92:1357
 Nitta R, Nishikawa T. Increased T-wave amplitude after accidental intravascular injection of lidocaine plus bupivacaine without epinephrine in sevoflurane-anesthetized child, 92:915
- Tanaka T, see Tokioka H
 Tang C-S, Fan S-Z, Chan C-C. Smoking status and body size increase carbon monoxide concentrations in the breathing circuit during low-flow anesthesia, 92:542
- Tang J
 White PF, Wender RH, Naruse R, Kariger R, Sloninsky A, Karlan MS, Uyeda RY, Karlan SR, Reichman C, Whetstone B. Fast-track office-based anesthesia: a comparison of propofol versus desflurane with antiemetic prophylaxis in spontaneously breathing patients, 92:95
 see Zhou TJ
- Tang Y-S, see Janelle GM
 Tanigawa K, Takeda T, Goto E, Tanaka K. The efficacy of esophageal detector devices in verifying tracheal tube placement: a randomized cross-over study of out-of-hospital cardiac arrest patients, 92:375
- Taniuchi S, see Inada T
 Tan P-H, Liu K, Peng C-H, Yang L-C, Lin C-R, Lu C-Y. The effect of dexamethasone on postoperative pain and emesis after intrathecal neostigmine, 92:228
- Tan PPC, see Li J-Y
- Tarabykin A, see Fredman B
 Tarnapolski A, see Eilig I
 Tarshis J, see Morgan PJ
 Taylor MD, see Blanshard HJ
 Teague BA, see Kain ZN
 Tempe DK. In search of a reliable awareness monitor (editorial), 92:801
- Tetzlaff JE
 Local and Regional Anesthesia; From the *Fundamentals of Anaesthesia and Acute Medicine Series* (book review), 92:1620
 see O'Hara JF Jr
- Thal GD. *Frontiers in Anesthesia; Volume 12, Number 1 of Problems in Anesthesia* (book review), 92:1078
- Thangathurai D, see Roffey P
 Thorén A, Elam M, Ricksten S-E. Jejunal mucosal perfusion is well maintained during mild hypothermic cardiopulmonary bypass in humans, 92:5
- Thys DM, see Vloka JD
 Tobias JD, see Berkenbosch JW
 Tobias MD, see Ochroch EA
 Tobo K, see Nagata E
 Todorov A, see Zwetsch G
 Tokioka H, Tanaka T, Ishizu T, Fukushima T, Iwaki T, Nakamura Y, Kosogabe Y. The effect of breath termination criterion on breathing patterns and the work of breathing during pressure support ventilation, 92:161
- Tolan MM, see Mortero RF
 Toller WG, see Gross ER
 Tomihisa T, see Kawasaki T
 Tomioka S, Takechi M, Ohshita N, Nakajo N. Propofol is not effective for hyperventilation syndrome, 92:781
- Tomiyasu S, see Hara T
 Tongier WK, see Coloma M
 Toriumi T, see Yokota K
 Torresani T, see Simma B
 Torres L-M, see Calderón E
 Torri G, see Casati A
 Tortosa JA, see Hernández-Palazón J
 Tousignant CP, see Gillham MJ
 Toyama A, see Fujioka S
 Toyooka H, see Fujii Y; Fukuda T
 Trachez MM, see Zapata-Sudo G
 Tramèr MR, see Choi PT-L; Walder B
 Traub RD, see Dexter F
 Trautner H, see Greim C-A
 Trigo C, see Fernández-Guisasola J
 Tröger U, see Martens-Lobenhoffer J
 Tsai S-K, see Yeh H-M
 Tsai T-Y, see Chen T-Y
 Tsai Y-C, Chang P-J, Jou I-M. Direct tramadol application on sciatic nerve inhibits spinal somatosensory evoked potentials in rats, 92:1547
- Tschernko EM, see Gruber EM
 Tseng C-C, see Chen T-Y
 Tsen LC. The mask avenger? (letter), 92:279
- Tsubo T, see Suzuki A
 Tsuda A, Shibata O, Saito M, Hashimoto S, Iwanaga S, Makita T, Sumikawa K. A dose-response study of anticholinesterase drugs on contractile and phosphatidylinositol responses of rat trachea, 92:100
- Tsueda K, see Mortero RF
 Tsujito T, see Yanagidate F
 Tuman KJ, see O'Connor CJ
 Tung A, Lynch JP, Mendelson WB. Prolonged sedation with propofol in the rat does not result in sleep deprivation, 92:1232
- Tweed WA, Dakin D. Explosive coughing after bolus fentanyl injection, 92:1442
- Tzeng J-I, see Ho S-T
 Tzoufi MJ, see Mentzelopoulos SD
- Uchida K, see Orii R
 Uemura A, see Fujii Y
 Ulrich B, see Heidegger T
 Urdaneta F, see Janelle GM
 Usui T, Saito S, Goto F. Arytenoid dislocation while using a McCoy laryngoscope, 92:1347
- Uyeda RY, see Tang J
- VadeBoncouer T, see Weinberg G
 Vahle-Hinz C, Detsch O, Siemers M, Kochs E, Bromm B. Local GABA_A receptor blockade reverses isoflurane's suppressive effects on thalamic neurons *in vivo*, 92:1578
- Vaisman J. Pelvic hematoma after an ilioinguinal nerve block for orchialgia, 92:1048
- Vakharia AS, see White PF
 Vakkuri A, see Jäntti V
 Vaknin Z, Manisterski Y, Ben-Abraham R, Efrati O, Lotan D, Barzilay Z, Paret G. Is endotracheal adrenaline deleterious because of the beta adrenergic effect?, 92:1408
- Vale PR, see Lathi KG
 Vallejo MC
 Adler LJ, Finegold H, Mandell GL, Ramanathan S. Periosteal entrapment of an epidural catheter in the intrathecal space, 92:1532
 Ramanathan S. Postdural puncture headache in obstetric patients

- (letter, reply), 92:1616
- Van Aken H, see Booke M; Brodner G
- van den Berg AA, see Neuvonen PT
- Vanderelst PE, see Singelyn FJ
- van der Woerd A, see Kerssens C
- Van Gessel E, see Culebras X
- Vatsgar TT, see Ræder JC
- Vela-Cantos F, see Leyvi G
- Venitz J, see Wahr JA
- Verghese ST, see Patel RI
- Vergnion M, Degesves S, Garcet L, Magotteaux V. Tramadol, an alternative to morphine for treating posttraumatic pain in the prehospital situation, 92:1543
- Vernon JC, see Groban L
- Veyckemans F, see Lavand'homme P; Singelyn FJ
- Vicenzi MN, see Mahla E
- Vig S, see Kaye AD
- Villa Y, see Berkenstadt H
- Virella CD, see Greilich PE
- Vissers KCP, see Vranken JH
- Vivien B, see David J-S
- Vizquel L, see Bernard J-M
- Vloka JD
- Hadžić A, April E, Thys DM. Anterior approach to the sciatic nerve block: the effects of leg rotation, 92:460
- Hadžić A, April E, Thys DM. The division of the sciatic nerve in the popliteal fossa: anatomical implications for popliteal nerve blockade, 92:215
- Voelckel WG
- Lurie KG, Zielinski T, McKnite S, Plaisance P, Wenzel V, Lindner KH. The effects of positive end-expiratory pressure during active compression decompression cardiopulmonary resuscitation with the inspiratory threshold valve, 92:967
- see Krismer AC
- Von Dossow V, Welte M, Zaune U, Martin E, Walter M, Rückert J, Kox WJ, Spies CD. Thoracic epidural anesthesia combined with general anesthesia: the preferred anesthetic technique for thoracic surgery, 92:848
- Von Eckardstein A, see Brodner G
- Vorwerk R, see Hering R
- Vranken JH, Vissers KCP, de Jongh R, Heylen R. Intraarticular sufentanil administration facilitates recovery after day-case knee arthroscopy, 92:625
- Vycudilik W, see Fitzgerald RD
- Wada H, see Higuchi H
- Wadhwa A, Clarke D, Goodchild CS, Young D. Large-dose oral dextromethorphan as an adjunct to patient-controlled analgesia with morphine after knee surgery, 92:448
- Wafai Y, see Nimmagadda U
- Wahr JA, Gerber M, Venitz J, Baliga N. Allosteric modification of oxygen delivery by hemoglobin, 92:615
- Waintrop C, see Basto ER
- Wajima Z, Yoshikawa T, Ogura A, Imanaga K, Shiga T, Inoue T, Ogawa R. The effects of diltiazem on hemodynamics and seizure duration during electroconvulsive therapy, 92:1327
- Wakamatsu H, see Matsumoto M
- Walcourt A, Scott RL, Nash HA. Blockage of one class of potassium channel alters the effectiveness of halothane in a brain circuit of *Drosophila*, 92:535
- Walder B
- Elia N, Henzi I, Romand JR, Tramèr MR. A lack of evidence of superiority of propofol versus midazolam for sedation in mechanically ventilated critically ill patients: a qualitative and quantitative systematic review, 92:975
- see Choi PT-L
- Wall MH, see Sherman SV
- Walter M, see Von Dossow V
- Wang J-J, see Ho S-T; Liu Y-H
- Wang L-K, see Chen T-Y
- Wang M-J, see Yeh H-M
- Wang S-M
- Kain ZN. Auricular acupuncture: a potential treatment for anxiety, 92:548
- see Kain ZN
- Wang Y-F, see Ma H-C
- Warltier DC, see Gross ER; May JA
- Warner DO, see Hanazaki M
- Warren JF, see Greilich PE
- Watanabe H, see Kanaya N
- Watanabe K, see Adachi YU
- Watcha MF
- Investigations of the bispectral index monitor in pediatric anesthesia: first things first, 92:805
- see Fazi L
- Watson R, Leslie K. Nerve blocks versus subcutaneous infiltration for stereotactic frame placement, 92:424
- Weatherwax K, see Gan TJ
- Weinberg G, VadeBoncouer T. Improved energetics may explain the favorable effect of insulin infusion on bupivacaine cardiotoxicity (letter), 92:1075
- Weisel RD, see Hynninen M
- Welch KB, see Samra SK
- Welsh J, see McNulty SE
- Welte M, see Von Dossow V
- Wender RH, see Tang J
- Wenzel V, see Krismer AC; Prengel AW; Voelckel WG
- Werkgartner G, see Mahla E
- Werner C, see Himmelseher S
- Whetstone B, see Tang J
- White HC, see May JA
- White PF
- Bovill JG. From SIVA to ISAP!—a logical evolution (editorial), 92:1
- Ghoname EA, Ahmed HE, Hamza MA, Craig WF, Vakharia AS. The effect of montage on the analgesic response to percutaneous neuromodulation therapy, 92:483
- Li S, Chiu JW. Electroanalgesia: its role in acute and chronic pain management, 92:505
- see Coloma M; Kadar AG; Tang J; Zárte E; Zhou TJ
- White WD, see Newman MF
- Whitten CW, see Sun Y
- Widmann C, see Lang-Lazdunski L
- Wiesner G, Hoerauf K, Schroegendorfer K, Sobczynski P, Harth M, Ruediger HW. High-level, but not low-level, occupational exposure to inhaled anesthetics is associated with genotoxicity in the micronucleus assay, 92:118
- Wiley J, see Alexander JM; Philip J
- Williams HK, see Ben-David B
- Williams JP, see Paiste J
- Williamson R. Unlighted stylet tracheal intubation (letter), 92:1355
- Wilson RC, see Robinson AP
- Wilson SIO, see Sun Y
- Wirtz S, see Booke M
- Wisser W, see Gruber EM
- Wlody DJ, see Dimaculangan DP
- Woelckel HJ, see Wohlfeil ER
- Wohlfeil ER, Woelckel HJ, Gottschall JL, Poole W. Increased carboxyhemoglobin from hemolysis mistaken as intraoperative desflurane breakdown, 92:1609
- Woletz J, see Sprung J
- Wolf T, see Hemmerling TM
- Wong C-H, see Li J-Y
- Wong DT, see Cheng DCH
- Wong ELY, see Ngan Kee WD

- Wong JK, Haas DA, Hu JW. Local anesthesia does not block mustard-oil-induced temporomandibular inflammation, 92:1035
- Wrigge H, see Hering R
- Wu S, see Zhang Y
- Xiao W, Deng X. The minimum alveolar concentration of enflurane for laryngeal mask airway extubation in deeply anesthetized children, 92:72
- Xi S, see Jahr JS
- Xu W, see Chen J
- Yamada Y, see Oriti R
- Yamaguchi H, see Satsumae T
- Yamaguchi T, see Nishiyama T
- Yamamoto F, see Ide T
- Yamamoto K, see Asai T
- Yamamoto S, see Satsumae T
- Yamashita M, see Yokoyama M
- Yamashita S, see Satsumae T
- Yamazaki K, see Masaki E
- Yanagidate F
- Dohi S, Hamaya Y, Tsujito T. Reexpansion pulmonary edema after thorascopic mediastinal tumor resection, 92:1416
- Hamaya Y, Dohi S. Vaginal indigo carmine-induced severe hypotension (letter), 92:556
- see Ma H-C
- Yanagi K, see Fukuda T
- Yang L-C, see Tan P-H
- Yaster M, Aram L
- Yasunaga T, see Satsumae T
- Yeh H-M, Chen L-K, Shyu M-K, Lin C-J, Sun W-Z, Wang M-J, Mok MS, Tsai S-K. The addition of morphine prolongs fentanyl-bupivacaine spinal analgesia for the relief of labor pain, 92:665
- Yeomans DC, see Pirec V
- Yli-Hankala A, see Jäntti V
- Yokota K, Fujii T, Kimura K, Toriumi T, Sari A. Life-threatening hypoxemic respiratory failure after repair of acute type A aortic dissection: successful treatment with venoarterial extracorporeal life support using a prosthetic graft attached to the right axillary artery, 92:872
- Yokoyama M
- Itano Y, Mizobuchi S, Nakatsuka H, Kaku R, Takashima T, Hirakawa M. The effects of epidural block on the distribution of lymphocyte subsets and natural-killer cell activity in patients with and without pain, 92:463
- Mizobuchi S, Nagano O, Fujii H, Yamashita M, Hirakawa M. The effects of epidural insertion site and surgical procedure on plasma lidocaine concentration, 92:470
- Yoo KY, Lee JU, Kim HS, Jeong SW. The effects of opioids on isolated human pregnant uterine muscles, 92:1006
- Yoshikawa T, see Wajima Z
- Yoshitani K, Kawaguchi M, Sugiyama N, Sugiyama M, Inoue S, Sakamoto T, Kitaguchi K, Furuya H. The association of high jugular bulb venous oxygen saturation with cognitive decline after hypothermic cardiopulmonary bypass, 92:1370
- Young D, see Wadhwa A
- Yuasa H, Higashizawa T, Koga Y. Spinal anesthesia in human T lymphotropic virus type I-associated myelopathy (letter), 92:1618
- Zalzal G, see Finkel JC
- Zapata-Sudo G, Trachez MM, Sudo RT, Nelson TE. Is comparative cardiotoxicity of S(-) and R(+) bupivacaine related to enantiomer-selective inhibition of L-type Ca^{2+} channels?, 92:496
- Zárate E, Mingus M, White PF, Chiu JW, Scuderi P, Loskota W, Daneshgari V. The use of transcutaneous acupoint electrical stimulation for preventing nausea and vomiting after laparoscopic surgery, 92:629
- Zaugg M, Lachat ML, Pfammatter T, Cathomas G, Schmid ER. Sudden respiratory arrest resulting from brainstem embolism in a patient undergoing endovascular abdominal aortic aneurysm repair, 92:335
- Zaune U, see Von Dossow V
- Zaytoun GM, see Aouad MT
- Zhang Y
- Stabernack C, Sonner J, Dutton R, Eger EI II. Both cerebral $GABA_A$ receptors and spinal $GABA_A$ receptors modulate the capacity of isoflurane to produce immobility, 92:1585
- Wu S, Eger EI II, Sonner JM. Neither $GABA_A$ nor strychnine-sensitive glycine receptors are the sole mediators of MAC for isoflurane, 92:123
- Zhou TJ, Tang J, White PF. Propacetamol versus ketorolac for treatment of acute postoperative pain after total hip or knee replacement, 92:1569
- Ziegler-Pithamitsis D, see Himmelseher S
- Zielinski T, see Voelckel WG
- Zimmerman AA, see Gruber EM
- Zimmermann P, see Greim C-A
- Zinserling J, see Hering R
- Zoeggeler A, see Goll V
- Zohar E, see Fredman B
- Zurakowski D, see Gruber EM
- Zwetsch G, Filipovic M, Skarvan K, Todorov A, Seeberger MD. Transient recurrent laryngeal nerve palsy after failed placement of a transesophageal echocardiographic probe in an anesthetized patient, 92:1422

Subject Index

- Abdominal surgery**, see Surgery
- Acadesine**, see Pharmacology
- Acetaminophen**, see Analgesics
- Activated clotting time**, see Measurement techniques
- Acupuncture**, see Anesthetic techniques
- Acute fatty liver of pregnancy**, see Complications
- Addiction**, alcohol or drug, patient recovering from, special issues for anesthesiologist, 92:1601
- Adenosine triphosphate**, and NMDA receptor antagonists, anesthetic interaction, rats, 92:134
- Adhesion molecules**, see Inflammation
- Adrenaline**, see Pharmacokinetics
- Adrenergic receptors**, see Receptors
- Adrenomedullin**, see Sympathetic nervous system
- Adverse effects**, explosive coughing after bolus fentanyl injection, 92:1442
- Age factors**
adults vs infants, response to heparin, effect of low preoperative antithrombin activity, cardiac surgery patients, 92:66
apolipoprotein E polymorphisms and age at first CABG, 92:824
calculating induction dose of propofol, 92:656
elderly
absence of reflex tachycardia after spinal anesthesia (letter), 92:1357
patients undergoing ambulatory colonoscopy, monitored anesthesia care, remifentanyl vs meperidine, 92:80
older anesthesiologists, issues of concern, 92:1487
- Air embolism**, see Embolism
- Air quality**, PACU, online monitoring by proton-transfer-reaction mass spectrometry, 92:389
- Airway**
difficult
fiberscope-aided tracheal intubation, two-person technique, 92:1611
ILMA vs awake fiberoptic intubation, 92:1342
patient with Coffin-Siris syndrome, 92:554
hoarseness, postoperative, caused by Ortner's cardiopharyngeal syndrome, 92:1071
LMA removal, enflurane MAC, deeply anesthetized children, 92:72
occlusion, respiratory distress induced by, effect of epidural anesthesia, isoflurane-anesthetized cats, 92:749
severe laryngospasm, patient with superior laryngeal nerve injury, 92:271
smooth muscle
dose-response study of anticholinesterase drugs, rats, 92:100
effects of ethanol on Ca^{2+} sensitivity, 92:767
- Airway pressure release ventilation**, see Ventilation
- Alcohol addiction**, patient recovering from, special issues for anesthesiologist, 92:1601
- Alfentanil**, see Analgesics
- Allergy**, severe anaphylactic reaction to cisatracurium, child, 92:648
- Alveolar hemorrhage**, see Complications
- Ambulatory anesthesia**, see Anesthesia
- Amino acids**, see Brain
- Analgesics**
acetaminophen (paracetamol)
hepatic failure after acetaminophen and sevoflurane exposure, child, 92:1446
plus codeine, vs oral ibuprofen, for analgesia after ambulatory surgery, 92:1470
prodrug, propacetamol, intravenous administration, effect on morphine consumption after spinal fusion surgery, 92:1473
prodrug, propacetamol, vs ketorolac, for treatment of acute postoperative pain after total hip or knee replacement, 92:1569
rectal, prophylactically administered, and postoperative opioid requirements, elective cleft palate repair, children, 92:907
vs metamizol, postoperative analgesia at home after ambulatory hand surgery, 92:347
vs tramadol, postoperative analgesia at home after ambulatory hand surgery, 92:347
alfentanil, acute tolerance to, and inflammatory pain, 92:1296
codeine, acetaminophen plus, vs ibuprofen, for analgesia after ambulatory surgery, 92:1470
fentanyl
added to prolonged epidural infusions of ropivacaine after colonic surgery, 92:1276
bupivacaine with, addition of morphine, spinal analgesia for labor pain, 92:665
bupivacaine with, vs ropivacaine with fentanyl, for continuous epidural labor analgesia, 92:1024
bupivacaine with, vs ropivacaine with fentanyl, for postoperative PCEA, 92:1024
effects on spontaneous uterine contractility, 92:1006
epidural, sparing effect, levobupivacaine for epidural analgesia in labor, 92:410
intranasal, effect on emergence characteristics after sevoflurane anesthesia, children undergoing surgery for bilateral myringotomy tube placement, 92:1164
ropivacaine with, vs bupivacaine with fentanyl, for postoperative PCEA, 92:1024
ibuprofen, vs codeine/acetaminophen, for analgesia after ambulatory surgery, 92:1470
ketoprofen, intravenous, in thyroid and parathyroid surgery, 92:1052
ketorolac
intravenous, effect on opioid requirement and pain after cesarean delivery, 92:1010
sevoflurane and, patients undergoing breast surgery, 92:1058
vs propacetamol, for treatment of acute postoperative pain after total hip or knee replacement, 92:1569
meperidine
effects on spontaneous uterine contractility, 92:1006
PCEA, after cesarean delivery, 92:1010
postoperative analgesic requirements, effect of preemptive use of dextromethorphan, 92:739
vs remifentanyl, for monitored anesthesia care, older patients undergoing ambulatory colonoscopy, 92:80
metamizol
vs paracetamol, postoperative analgesia at home after ambulatory hand surgery, 92:347

Analgesics—continued

- vs tramadol, postoperative analgesia at home after ambulatory hand surgery, 92:347
- morphine
 - addition to fentanyl-bupivacaine spinal analgesia for labor pain, 92:665
 - effects on spontaneous uterine contractility, 92:1006
 - enhancement of myofilament Ca^{2+} sensitivity, intact guinea pig hearts, 92:602
 - epidural, PONV associated with, preventing, use of dexamethasone, dose-ranging study, 92:745
 - intraarticular, preemptive analgesic effect, ambulatory arthroscopic knee surgery, 92:923
 - intrathecal, vs intrathecal sufentanil, intrathecal sufentanil-morphine, and IV morphine PCA, for postthoracotomy pain, 92:31
 - IVPCA, vs intrathecal morphine, sufentanil, and their combination, for postthoracotomy pain, 92:31
 - N-type calcium channel blockers and, combined effects on Aδ vs C fiber mediated nociception, 92:239
 - PCA, after knee surgery, large-dose oral dextromethorphan as adjunct, 92:448
 - PCA, after spinal fusion surgery, effect of intravenous administration of propacetamol, 92:1473
 - PCA, after spinal fusion surgery, effects of intravenous administration of propacetamol, 92:1473
 - requirements, after thyroid and parathyroid surgery, and intravenous ketoprofen, 92:1052
 - sufentanil-morphine, intrathecal, vs intrathecal morphine, sufentanil, and IV morphine PCA, for postthoracotomy pain, 92:31
 - sustained-release, response of neuropathic pain and pain in complex regional pain syndrome I, patients pretreated with spinal cord stimulation, 92:488
 - tolerance, development, and modulations of spinal serotonin activity, 92:1563
 - vs sameridine and placebo, effects on hypercarbic ventilatory response, 92:529

- vs tramadol, treating posttraumatic pain in prehospital situation, 92:1543
- nefopam, analgesia, for extracorporeal shock wave lithotripsy (letter), 92:557
- remifentanil
 - as alternative to contraindicated neuraxial analgesia, parturients (letter), 92:1355
 - as labor analgesic, preliminary investigation (letter), 92:1358
 - cardiac surgery (letter), 92:557
 - effect on heat pain threshold, volunteers, 92:369
 - esmolol as alternative to, during desflurane anesthesia for fast-track outpatient gynecologic laparoscopic surgery, 92:352
 - for severe postoperative pain, comparison of two constant-dose continuous infusions, 92:715
 - vs meperidine, for monitored anesthesia care, older patients undergoing ambulatory colonoscopy, 92:80
- sameridine, vs morphine and placebo, effects on hypercarbic ventilatory response, 92:529
- sufentanil
 - effects on spontaneous uterine contractility, 92:1006
 - epidural sufentanil-ropivacaine and labor analgesia, 92:180
 - for PCEA during labor, 92:184
 - intraarticular, after day-case knee arthroscopy, 92:625
 - intrathecal, continuous, for postoperative analgesia, 92:244
 - intrathecal, vs intrathecal morphine, intrathecal sufentanil-morphine, and IV morphine PCA, for postthoracotomy pain, 92:31
 - morphine, intrathecal, vs intrathecal morphine, sufentanil, and IV morphine PCA, for postthoracotomy pain, 92:31
- tramadol
 - added to lidocaine, for intravenous regional anesthesia, 92:209
 - analgesic efficacy, impairment by concurrent administration of ondansetron, 92:1319
 - local anesthetic effects, 92:1547
 - vs metamizol, postoperative analgesia at home after ambulatory hand surgery, 92:347

- vs morphine, treating posttraumatic pain in prehospital situation, 92:1543
- vs paracetamol, postoperative analgesia at home after ambulatory hand surgery, 92:347
- urapidil, treatment of postanesthetic shivering, 92:257

Anaphylaxis

- intraoperative, postoperative management (letter), 92:559
- severe anaphylactic reaction to cisatracurium, child, 92:648

Anatomy

- brachial plexus, axillary part, ultrasonographic findings, 92:1271
- popliteal fossa, division of sciatic nerve, 92:215

Anesthesia

- ambulatory
 - anorectal surgery, dexamethasone, 92:85
 - awake craniotomy for removal of tumor, 92:89
 - effect of personality on midazolam pharmacokinetics, 92:621
 - esmolol vs remifentanil during desflurane anesthesia, 92:352
 - fast-tracking children, ambulatory surgery, 92:918
 - home postoperative analgesia, 92:347
 - intraarticular sufentanil in knee arthroscopy, 92:625
 - knee arthroscopy (letter), 92:556
 - oral ibuprofen vs codeine/acetaminophen, 92:1470
 - preemptive intraarticular morphine, 92:923
 - propofol and ketamine coadministration, 92:1465
 - propofol vs desflurane-antiemetics, office-based anesthesia, 92:95
 - rapacurium and ECT, 92:1171
 - remifentanil, monitored anesthesia care, colonoscopy, 92:80
 - transcutaneous acupoint stimulation and PONV, 92:629
- cardiovascular
 - acadesine and reperfusion lung injury, 92:590
 - adrenomedullin and jugular venous oxygen saturation, 92:837
 - allosteric modification of oxygen delivery by hemoglobin, 92:615
 - angiotensin II antagonists and anesthetic induction, 92:26
 - aortic dissection and hypoxemic respiratory failure, 92:872

- apolipoprotein E and age at first CABG, 92:824
- aprotinin dose management, 92:830
- basic TEE performance, 92:1103
- beta adrenergic effect of endotracheal adrenaline, 92:1408
- catheter-induced tricuspid regurgitation, 92:1117
- cerebral embolism in aortic stent-grafts, 92:335
- cerebral oxygenation and cognitive decline, 92:1370
- circulating blood volume and hemoglobin-based oxygen carriers, 92:609
- clinical onset of heparin, 92:1391
- constrictive pericarditis, 92:1424
- cooximetry interference by blood substitutes, 92:863
- debris and cell salvage bowls, 92:1137
- diaspirin cross-linked hemoglobin, 92:44
- diaspirin cross-linked hemoglobin and vascular tone, 92:324
- diastolic systolic function and cardiac surgery, 92:291
- effect of insulin on post-CPB arrhythmias, 92:810
- efficacy of intrathecal opioid after thoracotomy, 92:31
- embolic events during hip arthroplasty, 92:49
- epidural anesthesia, one-lung ventilation, and thoracic surgery, 92:848
- ethanol and stunned myocardium, 92:299
- explicit intraoperative recall at BIS of 47, 92:808
- fast-tracking after CABG (editorial), 92:1081
- fat elimination from autologous blood, 92:341
- gender does not influence ϵ -ACA concentration, 92:1384
- gene therapy for inoperable coronary disease, 92:19
- hematocrit, hypothermia, and neurological outcome, 92:329
- hemodynamic changes in aortic valve area, 92:584
- hemoglobin-based oxygen carrier during abdominal aneurysm repair, 92:1413
- heparin-induced thrombocytopenia and lepirudin, 92:344
- hydroxyethyl starch and O blood group, 92:1396
- hydroxyethyl starches and platelet function, 92:1402
- inhibition of Ca^{2+} transients by morphine, 92:602
- intraoperative recall at BIS 47, 92:808
- intravascular volume replacement and coagulation, 92:565
- ischemia monitoring with impedance, 92:12
- jejunal mucosal perfusion during bypass, 92:5
- maximal activated clotting time vs celite-activated clotting time, 92:578
- nitric oxide and coagulation, 92:320
- NO and one-lung ventilation, 92:842
- PentaLyte resuscitation, heparinoids, and thrombotic mediators, 92:314
- predicted risk of fatal heart failure, 92:596
- prediction of postoperative hypovolemia, 92:1146
- preischemic phosphodiesterase type III inhibition, 92:1377
- protamine and myocardium contractility, 92:817
- pulmonary arterial pressure and TEE, 92:1364
- pulmonary blood flow and PETCO₂, 92:306
- pulmonary embolism and aprotinin, 92:1418
- recurrent laryngeal nerve palsy and TEE probe, 92:1422
- reexpansion pulmonary edema and thoracoscopy, 92:1416
- remifentanyl and fentanyl in fast-track CABG, 92:1094
- remifentanyl vs fentanyl in CABG, 92:1084
- resuscitation, local anesthetic cardiotoxicity, 92:37
- safety of TEE, 92:1126
- search for reliable awareness monitor (editorial), 92:801
- sevoflurane and mitochondrial K_{ATP} channels, 92:1139
- TEE and esophageal injury, 92:870
- TEE and thrombosis of aortic valve prosthesis, 92:1123
- TEE imaging of differential perfusion cannula, 92:338
- TEG and major abdominal surgery, 92:572
- tranexamic acid levels during CPB, 92:1131
- transnasal TEE and foramen ovale patency, 92:1111
- Voluven, novel hydroxyethyl starch, 92:855
- critical care and trauma
 - acute lung injury and prone positioning, 92:1226
 - adhesion molecules and TGF- β 1 in ventilator-induced lung injury, 92:428
 - breath termination criterion during pressure support ventilation, 92:161
 - effects of propofol sedation on sleep, 92:1232
 - endobronchial lidocaine administration, 92:1505
 - endogenous vs exogenous vasopressin during CPR, 92:1499
 - factor- β 1 in ventilator-induced lung injury, 92:437
 - gas exchange during CPAP and airway pressure release ventilation, 92:950
 - Gram stain of mini-bronchoalveolar lavage fluid, 92:442
 - Lazarus syndrome, 92:688
 - olprinone of endotoxemia, 92:676
 - ONO1714 and diaphragm in sepsis, 92:959
 - paradoxical fat embolism, 92:688
 - PEEP during active compression decompression CPR, 92:967
 - propofol vs midazolam in ICU, 92:975
 - selective iNOS inhibition, septic rats, 92:681
 - sepsis, neutrophils, and IV anesthetics, 92:437
- electroconvulsive therapy (ECT)
 - hemodynamics and seizure duration, effects of diltiazem, 92:1327
 - rapacuronium as alternative to succinylcholine, 92:1171
- neurosurgical
 - anesthesia for stereotactic frame placement, 92:424
 - 17 β -estradiol and cerebral ischemia, 92:1520
 - continuous positive airway pressure and cerebral blood flow, 92:415
 - dexmedetomidine and awake craniotomy, 92:1251
 - direct current auditory evoked potentials, perioperative period, 92:154
 - effects of xenon on intracranial pressure, 92:1245
 - fentanyl/remifentanyl for SSEPs monitoring, 92:1510
 - FK506 and spinal cord ischemia, 92:1237
 - hemodynamics during scalp infiltration, 92:1516

Anesthesia—continued

- ischemic tolerance and spinal cord, 92:418
- monitoring in Germany, 1997 vs 1991, 92:166
- myelopathy and laminectomy, 92:1254
- outcome from head trauma plus hemorrhage, 92:669
- peripheral venous pressure, 92:172
- stroke volume variation, fluids, and craniotomy, 92:984
- vasomotion in subarachnoid hemorrhage, 92:990
- obstetric
 - acute fatty liver of pregnancy, 92:1268
 - bupivacaine vs ropivacaine and epidural analgesia, 92:1261
 - effect of magnesium on coagulation and platelet function, 92:1257
 - epidural anesthesia and polycythemia, 92:1535
 - epidural entrapment, 92:1532
 - epidural sufentanil-ropivacaine and labor analgesia, 92:180
 - intense labor pain and obstructed labor, 92:1524
 - intravascular volume before obstetric spinal anesthesia, 92:997
 - ketorolac after cesarean delivery, 92:1010
 - Larsen syndrome and short stature, 92:1266
 - levobupivacaine sparing by fentanyl, 92:410
 - opioids and uterine contractility *in vitro*, 92:1006
 - postdural puncture headache (letter), 92:1616
 - ropivacaine epinephrine test dose, 92:1529
 - spinal morphine prolongs labor analgesia, 92:665
 - sufentanil for PCEA, 92:184
 - transient neurologic symptoms, after spinal anesthesia with lidocaine, 92:405
 - transient neurologic symptoms, lidocaine-induced, postcesarean delivery, 92:401
- pain medicine
 - α -adrenergic antagonists and thermal hyperalgesia, 92:1558
 - pneumocephalus after cervical epidural steroid injection, 92:1576
 - propacetamol vs ketorolac after joint replacement, 92:1569
 - spinal serotonin and morphine tolerance, 92:1563
- pediatric

- allergy to cisatracurium, 92:648
- anesthetic interference and lasers, 92:1444
- antithrombin activity and heparin anticoagulation, 92:66
- bilateral vocal cord palsy, 92:352
- BIS and sevoflurane anesthesia, 92:877
- BIS monitoring (editorial), 92:805
- carnitine palmitoyltransferase I coma, 92:646
- caudal anesthesia, awake neonates, 92:1439
- clonidine and midazolam in tonsillectomy, 92:56
- CO₂ monitoring, end-tidal vs transcutaneous, 92:1427
- comparison of two ventilators, infants, 92:76
- continuous fascia iliaca block, 92:1159
- CSF leak with caudal blood patch, 92:1169
- enflurane MAC for LMA removal, 92:72
- failure of chain-link mechanism, 92:913
- fast-tracking after ambulatory surgery, 92:918
- fentanyl-induced coughing, 92:1442
- hepatic failure after acetaminophen and sevoflurane, 92:1446
- hypertonic saline and antidiuretic hormone, severe head injury, 92:641
- influence of sevoflurane and halothane on cerebral blood flow velocity, 92:891
- intranasal fentanyl and emergence, 92:1164
- IV dexamethasone for adenotonsillectomy, 92:636
- local anesthetics and T-wave increase, 92:913
- perioperative analgesia and acetaminophen, 92:907
- proper probe positioning for infants (letter), 92:1076
- sensory stimuli, preoperative anxiety, 92:897
- sevoflurane and congenital heart disease, 92:1152
- S(+)-ketamine for rectal premedication, 92:62
- Sonoclot analysis, 92:904
- stress response in cardiac anesthesia, 92:882
- thoracic, 92:1449
- tunneled epidural catheters, 92:1432
- regional
 - local anesthetic effects of tramadol, 92:1547

- superficial cervical blocks after thyroidectomy, 92:1538
- thoracic epidural for mastectomy, 92:1552
- tramadol after trauma in prehospital situation, 92:1543
- regional anesthesia and pain medicine
 - brachial plexus block at humeral canal, 92:194
 - breakage of epidural catheters, 92:246
 - bupivacaine enantiomers on Ca²⁺ channels, 92:496
 - Ca⁺⁺ channel blockers and morphine, 92:239
 - carbamazepine and morphine in neuropathic pain, 92:488
 - cardiac arrest during spinal anesthesia, 92:252
 - cardiac output after spinal anesthesia, 92:710
 - cervical block, ropivacaine vs bupivacaine, 92:720
 - chronic pain suppression with dietary soy, 92:1029
 - clonidine and glutamate antagonists, 92:725
 - clonidine in brachial plexus block, 92:199
 - continuous vs PCA femoral nerve sheath block, 92:455
 - dexamethasone and emesis, 92:745
 - dexamethasone and spinal neostigmine, 92:228
 - dextromethorphan for pain relief after knee surgery, 92:448
 - diffuse brachial plexopathy after interscalene blockade, patient receiving cisplatin, 92:249
 - division of sciatic nerve in popliteal fossa, 92:215
 - effect of epidural block on lymphocyte subsets, 92:463
 - effect of percutaneous neuromodulation therapy montage on analgesic response, 92:483
 - EP₁ antagonist and postoperative pain, 92:233
 - epidural anesthesia and respiratory distress, 92:749
 - epidural infusions of ropivacaine, 92:1276
 - epidural S(+)-ketamine decreases postoperative pain, 92:1290
 - glucocorticoid inhibition of hyperalgesia, 92:476
 - inflammatory pain and acute tolerance to alfentanil, 92:1296
 - inhibition of synovial plasma extravasation, 92:1301

- interaction of clonidine and lidocaine, 92:733
- intrathecal sufentanil analgesia, 92:244
- IV regional anesthesia with tramadol, 92:209
- ketamine and tourniquet hypertension, 92:1286
- lysophosphatidic acid and neutrophils, 92:1041
- microfabricated microneedles, 92:502
- neurogenic temporomandibular inflammation, 92:1035
- ondansetron and tramadol consumption, 92:1319
- patient-controlled interscalene anesthesia and shoulder surgery, 92:218
- PCA wound instillation, 92:189
- pelvic hematoma after nerve block, 92:1048
- plasma lidocaine concentration during surgery, 92:470
- preemptive dextromethorphan, 92:739
- remifentanyl infusion for severe postoperative pain, 92:715
- reversible paraplegia and epidural analgesia, 92:1326
- ropivacaine or bupivacaine for peripheral nerve blocks, 92:205
- ropivacaine vs bupivacaine for PCEA, 92:1024
- sciatic nerve block and leg rotation, 92:460
- small-dose spinal anesthesia and TURP, 92:1020
- spino-subcutaneous fistula treated with epidural blood patch, 92:1050
- supraspinal and spinal action of cholinomimetics, 92:1307
- survey of patients after PCA, 92:224
- thoracic epidural anesthesia and breathing, 92:1015
- thoracic epidural anesthesia for bilateral reduction mammoplasty, 92:514
- ultrasonic brachial plexus findings, 92:1271
- use of electroanalgesia for pain management, 92:505
- Anesthesia & Analgesia**, see Publications
- Anesthesiologists**
- adequacy of basic TEE performance, 92:1103
- aging, issues of concern, 92:1487
- masks, use of, during neuraxial block placement (letters), 92:279
- occupational exposure, air quality, PACU, online monitoring by proton-transfer-reaction mass spectrometry, 92:389
- Anesthetic machines**, see Equipment
- Anesthetics, gases**
- nitrous oxide (N_2O)
- enhancement of ketamine-induced c-Fos expression, posterior cingulate and retrosplenial cortices, rats, 92:362
- filled endotracheal tube cuff, deflationary phenomenon after cessation of N_2O administration, 92:145
- N_2O gas-barrier properties, Profile Soft-Seal Cuff, 92:140
- occupational exposure, high-level vs low-level, association with genotoxicity, micronucleus assay, 92:118
- sevoflurane in, children anesthetized with, effect of BIS monitoring on anesthetic use and recovery, 92:877
- sevoflurane- N_2O , vs thiopental/succinylcholine, endotracheal intubation, adults, 92:523
- xenon
- inhalation, effects on pial vessel diameter and intracranial pressure, rabbits, 92:1245
- inhibition of ketamine-induced c-Fos expression, posterior cingulate and retrosplenial cortices, rats, 92:362
- Anesthetics, intravenous**
- dexamethasone, effect on PONV after tonsillectomy, 92:636
- dexmedetomidine, infusion, for awake craniotomy, 92:1251
- fentanyl
- bolus injection, followed by explosive coughing, 92:1442
- bupivacaine spinal analgesia for relief of labor pain, addition of morphine, 92:665
- fast-tracking after CABG (editorial), 92:1081
- remifentanyl and, efficacy and resource utilization in fast-track CABG surgery, 92:1094
- stress response in infants undergoing cardiac surgery, 92:882
- vs remifentanyl, for intraoperative monitoring of SSEPs, 92:1510
- vs remifentanyl, when combined with isoflurane/propofol, for early extubation, CABG, 92:1084
- ketamine
- induced c-Fos expression, posterior cingulate and retrosplenial cortices, effects of xenon and N_2O , rats, 92:362
- modulation of NMDA receptor function, 92:1173
- modulation of NMDA receptor function, interactions with volatile anesthetics, 92:1182
- S(+)-ketamine, for rectal premedication, children, 92:62
- small-dose, effects on propofol sedation, 92:1465
- small-dose, preoperative, prevention of tourniquet-induced arterial pressure increase, orthopedic patients under general anesthesia, 92:1286
- small-dose S(+)-, applied with ropivacaine in epidural anesthesia, reduction of postoperative pain, total knee arthroplasty, 92:1290
- methohexital, direct current auditory evoked potentials, perioperative period, 92:154
- midazolam
- effectiveness for hyperventilation syndrome, 92:781
- stress response in infants undergoing cardiac surgery, 92:882
- vs propofol, effects on neutrophil hydrogen peroxide production and adhesion molecule expression, abdominal sepsis, rats, 92:437
- vs propofol, for sedation, mechanically ventilated critically ill patients, 92:975
- propofol
- adequate level of anesthesia, auditory information processing during, monitored by BIS, 92:1210
- block of muscle sodium channels, voltage-dependent manner, 92:1192
- direct current auditory evoked potentials, perioperative period, 92:154
- effects in area postrema, rats, 92:934
- induction dose, determinants, 92:656
- ineffectiveness for hyperventilation syndrome, 92:781
- prolonged sedation with, effects on sleep, rats, 92:1232
- sedation, effects of small-dose ketamine, 92:1465
- vs desflurane, with antiemetic prophylaxis, fast-track office-based anesthesia, 92:95
- vs midazolam, effects on neutrophil hydrogen peroxide production and adhesion molecule expression, abdominal sepsis, rats, 92:437

Anesthetics, intravenous—continued

vs midazolam, for sedation,
mechanically ventilated critically ill
patients, 92:975

remifentanyl

fast-tracking after CABG (editorial),
92:1081

fentanyl and, efficacy and resource
utilization in fast-track CABG
surgery, 92:1094

vs fentanyl, for intraoperative
monitoring of SSEPs, 92:1510

vs fentanyl, when combined with
isoflurane/propofol, for early
extubation, CABG, 92:1084

thiopental

inhibition of human $\alpha 7$ nicotinic
acetylcholine receptor, 92:930
/succinylcholine, vs sevoflurane- N_2O ,
endotracheal intubation, adults,
92:523

Anesthetics, local

benzocaine, factors influencing
Hurracaine delivery, 92:379

bupivacaine

0.5%, vs 0.5% ropivacaine, minimum
local anesthetic volume blocking
femoral nerve in 50% of cases,
92:205

0.15%, vs ropivacaine 0.2%, PCIA after
major open shoulder surgery, effects
on hand motor function, 92:218

0.25%, thoracic epidural analgesia,
effects on ventilatory mechanics,
patients with severe chronic
obstructive pulmonary disease,
92:1015

0.125%, sufentanil added to, dose-
range effects on quality of PCEA
during labor, 92:184

0.0625%, with fentanyl, vs 0.1%
ropivacaine with fentanyl, for
continuous epidural labor analgesia,
92:1261

cardiotoxicity, favorable effect of
insulin infusion (letter), 92:37

clonidine combined with, brachial
plexus block, postoperative
analgesia and hemodynamic
changes, 92:199

continuous fascia iliaca compartment
block, children, 92:1159

fentanyl-bupivacaine spinal analgesia
for relief of labor pain, addition of
morphine, 92:665

incremental overdosage, cardiac
resuscitation after, anesthetized
dogs, 92:37

lidocaine-bupivacaine mixture, T-
wave increase after accidental
intravascular injection, 92:915

neurogenic temporomandibular
inflammation, 92:1035

PCA wound instillation, 92:189

S(-) and R(+) bupivacaine,
comparative cardiotoxicity, and
enantiomer-selective inhibition of
L-type Ca^{2+} channels, 92:496

vs ropivacaine, ambulatory labor
epidural analgesia, potencies and
probabilities (letter), 92:278

vs ropivacaine, for cervical plexus
block, 92:720

with fentanyl, vs ropivacaine with
fentanyl, for postoperative PCEA,
92:1024

wound instillation, via electronic PCA
device and double-catheter system,
failure to decrease postoperative
pain or opioid requirements after
major abdominal surgery, 92:189

levobupivacaine

for epidural analgesia in labor,
sparing effect of epidural fentanyl,
92:410

incremental overdosage, cardiac
resuscitation after, anesthetized
dogs, 92:37

lidocaine

attenuation of lysophosphatidic acid-
induced priming in neutrophils,
92:1041

-bupivacaine mixture, T-wave increase
after accidental intravascular
injection, 92:915

effects on NO production from
activated murine macrophage cell
line, 92:128

endobronchial administration,
endotracheal tube vs LMA, 92:1505

incremental overdosage, cardiac
resuscitation after, anesthetized
dogs, 92:37

-induced transient neurologic
symptoms, postcesarean section,
92:401

intracuff, use during general
anesthesia (letter), 92:1075

neurogenic temporomandibular
inflammation, 92:1035

neurotoxicity, obstetric patients
(editorial), 92:287

plasma concentration, effects of
epidural insertion site and surgical
procedure, 92:470

spinal anesthesia with, transient
neurologic symptoms after, obstetric
patients, 92:405

spinal clonidine and, antinociceptive
interaction between, rat formalin
test, isobolographic analysis, 92:733
tramadol added to, for intravenous
regional anesthesia, 92:209
with epinephrine, scalp infiltration,
cardiovascular responses,
craniotomy, 92:1516

ropivacaine

0.1%, with fentanyl, vs 0.0625%
bupivacaine with fentanyl, for
continuous epidural labor analgesia,
92:1261

0.2%, vs bupivacaine 0.15%, PCIA
after major open shoulder surgery,
effects on hand motor function,
92:218

0.5%, vs 0.5% bupivacaine, minimum
local anesthetic volume blocking
femoral nerve in 50% of cases,
92:205

attenuation of lysophosphatidic acid-
induced priming in neutrophils,
92:1041

epidural S(+)-ketamine decreases
postoperative pain, 92:1290

epidural sufentanil-ropivacaine and
labor analgesia, 92:180

incremental overdosage, cardiac
resuscitation after, anesthetized
dogs, 92:37

prolonged epidural infusions, after
colonic surgery, impact of adding
fentanyl, 92:1276

thoracic epidural for mastectomy,
92:1552

vs bupivacaine, ambulatory labor
epidural analgesia, potencies and
probabilities (letter), 92:278

vs bupivacaine, for cervical plexus
block, 92:720

with epinephrine, as epidural test
dose in parturients, limitations,
92:1529

with fentanyl, vs bupivacaine with
fentanyl, for postoperative PCEA,
92:1024

sameridine, vs morphine and placebo,
effects on hypercarbic ventilatory
response, 92:529

tetracaine

attenuation of lysophosphatidic acid-
induced priming in neutrophils,
92:1041

small-dose spinal anesthesia for
transurethral prostatectomy, 92:1020

Anesthetics, volatile

desflurane

- anesthesia, fast-track outpatient
 - gynecologic laparoscopic surgery, esmolol as alternative to remifentanyl, 92:352
- breakdown, intraoperative, increased carboxyhemoglobin from hemolysis mistaken for, 92:1609
- carbon monoxide during anesthesia, 92:542
- NMDA receptor signaling and, 92:1182
- vs sevoflurane, cerebral hemodynamic responses (letter), 92:282
- with antiemetic prophylaxis, vs propofol, fast-track office-based anesthesia, 92:95
- enflurane, MAC for LMA extubation, deeply anesthetized children, 92:72
- halothane
 - cerebral hyperemic response, augmentation by endotoxin (letter), 92:1075
 - effectiveness in brain circuit of *Drosophila*, effects of blockage of one class of potassium channel, 92:535
 - vs sevoflurane, effects on cerebral blood flow velocity, children, 92:891
 - vs sevoflurane, safety and efficacy, infants and children with congenital heart disease, 92:1152
- isoflurane
 - carbon monoxide during anesthesia, 92:542
 - effects on renal function, patients with stable moderate renal insufficiency, 92:650
 - immobilizing action, modulation by cerebral and spinal GABA_A receptors, 92:1585
 - MAC, mediators, GABA_A and strychnine-sensitive glycine receptors, 92:123
 - NMDA receptor signaling and, 92:1182
 - suppressive effects on thalamic neurons *in vivo*, reversal by local GABA_A receptor blockade, 92:1578
- occupational exposure, high-level vs low-level, association with genotoxicity, micronucleus assay, 92:118
- sevoflurane
 - ATP and NMDA receptor interactions, 92:134
 - epileptogenic property, patients without epilepsy (letter), 92:1359

- hepatic failure after acetaminophen and sevoflurane exposure, child, 92:1446
- in N₂O, children anesthetized with, effect of BIS monitoring on anesthetic use and recovery, 92:877
- ketorolac and, patients undergoing breast surgery, 92:1058
- low-flow, effects on renal function, patients with stable moderate renal insufficiency, 92:650
- NMDA receptor signaling and, 92:1182
- N₂O, vs thiopental/succinylcholine, endotracheal intubation, adults, 92:523
- protection of stunned myocardium through activation of mitochondrial ATP-sensitive potassium channels, 92:1139
- vs desflurane, cerebral hemodynamic responses (letter), 92:282
- vs halothane, effects on cerebral blood flow velocity, children, 92:891
- vs halothane, safety and efficacy, infants and children with congenital heart disease, 92:1152

Anesthetic techniques

- acupuncture, auricular, potential treatment for anxiety, 92:548
- brachial plexus block
 - at humeral canal, using neurostimulator, evaluation, 92:194
- axillary approach, ultrasonic brachial plexus findings, 92:1271
- effect of clonidine combined with long-acting local anesthetic, 92:199
- interscalene, difficult insertion of catheter (letter), 92:1074
- caudal
 - local anesthetics and T-wave, 92:915
 - single-dose, for major intraabdominal operations, high-risk infants, 92:1439
- cervical plexus block
 - bilateral superficial, prevention of postoperative pain, thyroid surgery, 92:1538
 - ropivacaine vs bupivacaine, 92:720
- chronic methylprednisolone infusion, inhibition of neuropathic hyperalgesia and spinal Fos expression, 92:476
- combined spinal-epidural analgesia, for relief of labor pain, 92:665
- conscious sedation, dexmedetomidine and awake craniotomy, 92:1251
- continuous infusion, remifentanyl, for severe postoperative pain,

- comparison of two constant-dose continuous infusions, 92:715
- continuous intrathecal sufentanil analgesia, 92:244
- electroanalgesia
 - percutaneous neuromodulation therapy, analgesic response, effect of montage, 92:483
 - role in acute and chronic pain management, 92:505
- endobronchial lidocaine, administration, endotracheal tube vs LMA, 92:1505
- epidural
 - analgesia, and reversible paraplegia, 92:1316
 - analgesia, labor, bupivacaine vs ropivacaine, 0.0625% bupivacaine with fentanyl vs 0.1% ropivacaine with fentanyl, 92:1261
 - analgesia, labor, bupivacaine vs ropivacaine, potencies and probabilities (letter), 92:278
 - analgesia, labor, levobupivacaine, sparing effect of epidural fentanyl, 92:410
 - analgesia, labor, sufentanil-ropivacaine, 92:180
 - analgesia, patient with paroxysmal nocturnal hemoglobinuria (letter), 92:1618
 - analgesia, prolonged, tunneled epidural catheters, pediatric patients, 92:1432
 - block, effects on distribution of lymphocyte subsets and natural-killer cell activity, patients with and without pain, 92:463
 - catheters, "blocked," another cause (letter), 92:1617
 - catheters, breakage, 92:246
 - catheters, periosteal entrapment in intrathecal space, 92:1532
 - catheters, tunneled, for prolonged analgesia, pediatric patients, 92:1432
 - continuous, insertion site, effects on plasma lidocaine concentration, 92:470
 - for cesarean section, patient with polycythemia rubra vera and preeclampsia, 92:1535
 - parturient with Larsen syndrome, 92:1266
 - pediatric thoracic anesthesia, 92:1449
 - ropivacaine epinephrine test dose, 92:1529
 - S(+)-ketamine decreases postoperative pain, 92:1290
- epidural blood patch

Anesthetic techniques—continued

- caudal approach, treatment of intractable PONV after placement of permanent intrathecal catheter, 92:1169
- successful treatment of spino-subcutaneous fistula after bone marrow harvest, 92:1050
- extended femoral nerve sheath block, after total hip arthroplasty, continuous vs PCA, 92:455
- fascia iliaca compartment block, continuous, in children, 92:1159
- fast-tracking
 - after CABG (editorial), 92:1081
 - CABG surgery, efficacy and resource utilization of remifentanyl and fentanyl, 92:1094
 - children after ambulatory surgery, 92:918
 - esmolol vs remifentanyl during desflurane anesthesia, 92:352
 - office-based anesthesia, propofol vs desflurane with antiemetic prophylaxis, 92:95
- femoral nerve block, minimum local anesthetic volume, 0.5% ropivacaine and 0.5% bupivacaine, 92:205
- general
 - combined with thoracic epidural anesthesia, for thoracic surgery, 92:848
 - ketamine and tourniquet hypertension, 92:1286
 - use of intracuff lidocaine during (letter), 92:1075
 - vs continuous thoracic epidural, for modified radical mastectomy, 92:1552
- ilioinguinal nerve block, for orchialgia, pelvic hematoma after, 92:1048
- interscalene block
 - brachial plexus catheter, difficult insertion (letter), 92:1074
 - diffuse brachial plexopathy, patient receiving cisplatin therapy, pharmacological double crush syndrome, 92:249
- intraarticular
 - morphine, preemptive analgesic effect, ambulatory arthroscopic knee surgery, 92:923
 - sufentanil, after day-case knee arthroscopy, 92:625
- intranasal fentanyl, effect on emergence characteristics after sevoflurane anesthesia, children undergoing surgery for bilateral myringotomy tube placement, 92:1164
- intravenous regional, tramadol added to lidocaine, 92:209
- low-flow anesthesia, carbon monoxide concentrations in breathing circuit, effects of smoking status and body size, 92:542
- monitored anesthesia care
 - awake craniotomy for removal of tumor, 92:89
 - effects of small-dose ketamine on propofol sedation, 92:1465
 - remifentanyl vs meperidine, older patients undergoing ambulatory colonoscopy, 92:80
- nerve blocks, vs subcutaneous infiltration, for stereotactic frame placement, 92:424
- neuraxial block
 - analgesia, contraindicated in parturients, remifentanyl as alternative (letter), 92:1355
 - placement, use of face masks during (letters), 92:279
- orbital block, use of hyaluronidase (letter), 92:560
- paravertebral block, characteristics (letter), 92:558
- patient-controlled analgesia (PCA)
 - electronic PCA device, and double-catheter system, bupivacaine wound instillation, failure to decrease postoperative pain or opioid requirements after major abdominal surgery, 91:189
- extended femoral nerve sheath block after total hip arthroplasty, 92:455
- IV morphine, vs intrathecal morphine, sufentanil, and their combination, for postthoracotomy pain, 92:31
- ondansetron and tramadol consumption, 92:1319
- propacetamol and morphine consumption after spinal fusion surgery, 92:1473
- propacetamol vs ketorolac after joint replacement, 92:1569
- prospective survey of patients after cessation, 92:224
- with morphine, after knee surgery, large-dose oral dextromethorphan as adjunct, 92:448
- wound instillation, after major abdominal surgery, 92:189
- wound instillation, local anesthetics (editorial), 92:3
- patient-controlled epidural analgesia (PCEA)
 - during labor, dose-range effects of sufentanil added to 0.125% bupivacaine, 92:184
 - meperidine, after cesarean delivery, 92:1010
 - postoperative, ropivacaine with fentanyl vs bupivacaine with fentanyl, 92:1024
- patient-controlled interscalene analgesia (PCIA)
 - after major open shoulder surgery, 92:218
- peripheral nerve block, eliciting paresthesias for (letter), 92:795
- preemptive analgesia
 - dextromethorphan, effect on postoperative analgesic requirements, 92:739
 - inhibition of synovial plasma extravasation, 92:1301
 - intraarticular morphine, arthroscopic knee surgery, 92:923
- rapid sequence induction, preoxygenation, tidal volume vs deep breaths, 92:1337
- rectal premedication, S(+)-ketamine, children, 92:62
- scalp infiltration, with epinephrine with or without lidocaine, cardiovascular responses, 92:1516
- sciatic nerve block
 - anterior approach, effects of leg rotation, 92:460
 - division of sciatic nerve in popliteal fossa, 92:215
- spinal
 - absence of reflex tachycardia after, elderly patients (letter), 92:1357
 - cardiac arrest during, common mechanisms and strategies for prevention, 92:252
 - cardiac output changes, effects of lactated Ringer's solution infusion, 92:710
 - in human T lymphotropic virus type I-associated myelopathy (letter), 92:1618
 - intravascular volume before obstetric spinal anesthesia, 92:997
 - lidocaine-induced transient neurological symptoms, postcesarean section, 92:401
 - lidocaine neurotoxicity, obstetric patients (editorial), 92:287
 - neostigmine, dexamethasone and, 92:228
 - small-dose, tetracaine, for transurethral prostatectomy, 92:1020

- with lidocaine, and transient neurologic symptoms, obstetric patients, 92:405
- spinal cord stimulation, patients pretreated with, response of neuropathic pain and pain in complex regional pain syndrome I to carbamazepine and sustained-release morphine, 92:488
- subcutaneous infiltration, vs nerve blocks, for stereotactic frame placement, 92:424
- thoracic epidural
 - analgesia, multimodal postoperative rehabilitation program, 92:1594
 - combined with general anesthesia, for thoracic surgery, 92:848
 - continuous, vs general anesthesia, for modified radical mastectomy, 92:1552
 - for bilateral reduction mammoplasty, patient with Klippel-Feil syndrome, 92:514
 - infusions of ropivacaine, prolonged, after colonic surgery, impact of adding fentanyl, 92:1276
 - respiratory distress induced by airway occlusion, isoflurane-anesthetized cats, 92:749
 - with bupivacaine 0.25%, effects on ventilatory mechanics, patients with severe chronic obstructive pulmonary disease, 92:1015
- three vital capacity breaths, endotracheal intubation, adults, 92:523
- topical, factors influencing Hurracaine delivery, 92:379
- transdermal drug delivery, lack of pain associated with microfabricated microneedles, 92:502
- tunneled epidural catheter, for prolonged analgesia, pediatric patients, 92:1432
- weight-related dose regimen, perioperative plasma aprotinin concentration, 92:830
- wound instillation, patient-controlled after major abdominal surgery, 92:189
- local anesthetics (editorial), 92:3
- Aneurysm repair**, see Surgery, abdominal
- Angiotensin II antagonists**, see Pharmacology
- Anorectal surgery**, see Surgery
- Antagonists, neuromuscular relaxants**
 - edrophonium, effects on contractile and phosphatidylinositol responses of trachea, dose-response study, rats, 92:100
 - neostigmine, effects on contractile and phosphatidylinositol responses of trachea, dose-response study, rats, 92:100
 - pyridostigmine, effects on contractile and phosphatidylinositol responses of trachea, dose-response study, rats, 92:100
- Anticholinesterase drugs**, see Pharmacology
- Antidiuretic hormone**, see Hormones
- Antiemetic techniques**, transcutaneous
 - acupoint electrical stimulation, for preventing PONV after laparoscopic surgery, 92:629
- Antifibrinolytics**
 - regional hemostatic status and blood requirements after total knee arthroplasty, 92:775
 - therapy, pulmonary thromboembolism and, during orthotopic liver transplantation (letter), 92:559
- Antithrombin activity**, see Coagulation
- Anxiety**, see Psychology
- Aorta**
 - ischemia-reperfusion injury, PentaLyte resuscitation, heparinoids, and thrombotic mediators, rabbits, 92:314
 - prosthetic valve, acute thrombosis, diagnosis by TEE, 92:1123
- Aortic valve area**, see Heart
- Apolipoprotein E**, see Cholesterol
- Aprotinin**, see Coagulation; Pharmacokinetics; Pharmacology
- APRV**, see Airway pressure release ventilation
- Area postrema**, see Brain
- Arrhythmias**, see Heart
- Arterial lines**, see Equipment
- Arthroscopy**, see Surgery, knee
- Arytenoid dislocation**, see Complications
- Atrial fibrillation**, see Heart
- Auditory evoked potentials**, see Monitoring
- Auditory information processing**, see Brain
- Awareness**, see Complications
- Axillary block**, see Anesthetic techniques
- Benzocaine**, see Anesthetics, local
- Bilateral vocal cord palsy**, see Complications
- Bispectral index (BIS)**, see Monitoring, electroencephalography
- Bleeding**, see Complications
- Blood**
 - autologous, fat elimination, 92:341
 - carboxyhemoglobin
 - cooximetry interference by hemoglobin-based blood substitutes, 92:863
 - increased, from hemolysis, mistaken for intraoperative desflurane breakdown, 92:1609
 - hematocrit, levels, varying, blood replacement with, neurological recovery, porcine model of profound hypothermic circulatory arrest, 92:329
 - hemoglobin
 - allosteric modification of oxygen delivery, 92:615
 - artificial, diaspirin cross-linked hemoglobin and vascular tone, 92:324
 - concentrations, after large-dose infusions of diaspirin cross-linked hemoglobin, 92:44
 - cooximetry interference by hemoglobin-based blood substitutes, 92:863
 - hemoglobin-based oxygen carrier
 - cooximetry interference, 92:863
 - measurement of circulating blood volume (rabbits), 92:609
 - successful use as primary blood substitute, abdominal aneurysm repair, 92:1413
 - hemolysis, increased carboxyhemoglobin mistaken for intraoperative desflurane breakdown, 92:1609
- loss
 - abdominal aneurysm repair, successful use of hemoglobin-based oxygen carrier as primary blood substitute, 92:1423
 - after total knee arthroplasty, regional hemostatic status and blood requirements with and without tranexamic acid or aprotinin, 92:775
 - diaspirin cross-linked hemoglobin, 92:44
 - in major orthopedic surgery, effects of Voluven vs HAES-steril, 92:855
 - methemoglobin, cooximetry interference by hemoglobin-based blood substitutes, 92:863
 - myeloproliferative disorder, patient with, epidural anesthesia for cesarean section, 92:1535
 - oxygen dissociation curve, allosteric modification of oxygen delivery by hemoglobin, 92:615
 - oxyhemoglobin, cooximetry interference by hemoglobin-based blood substitutes, 92:863

Blood—continued

- replacement, with varying hematocrit levels, neurological recovery, porcine model of profound hypothermic circulatory arrest, 92:329
- transfusion
 - autotransfusion techniques, cell salvage bowls, partly filled, debris elimination, 92:1137
 - diaspirin cross-linked hemoglobin, 92:44
 - fat elimination from autologous blood, 92:341
 - hydroxyethyl starch and O blood group, 92:1396
 - volume, circulating, measuring, use of hemoglobin-based oxygen carrier (rabbits), 92:609

Blood filters, see Equipment**Blood flow**, see Brain; Liver; Lung**Blood gas analysis**, see Measurement techniques**Blood groups**, O blood group, hemostatic changes in patients receiving hydroxyethyl starch, 92:1396**Blood pressure**

- decrease, caused by endotracheal adrenaline, mechanism, 92:1408
- hypertension
 - during ECT, effects of diltiazem, 92:1327
 - hemodynamics during scalp infiltration, 92:1516
 - treatment with angiotensin II antagonists, and anesthetic induction, 92:26
- hypotension
 - caused by sepsis, prevention by selective iNOS inhibition, rats, 92:681
 - hemodynamics during scalp infiltration, 92:1516
 - hypovolemic, after radical surgery for esophageal cancer, prediction, 92:1146
 - small-dose tetracaine spinal anesthesia for transurethral prostatectomy, 92:1020

Body size, see Weight**Book and multimedia reviews**

- Airway Management; Volume 38, Number 3, 92:798
- Anesthesia and Renal Considerations; Volume 18, Number 4 (December 2000) of *Anesthesiology Clinics of North America*, 92:1360
- Atlas of Pain Management Injection Techniques, 92:284

Awareness During Anesthesia, 92:1079

Basics of Anesthesia, 4th Edition, 92:1360

Current Issues in Anesthesiology;

- Volume 38, Number 4 of *International Anesthesiology Clinics*, 92:1620

Frontiers in Anesthesia; Volume 12, Number 1 of *Problems in Anesthesia*, 92:1078

Hypertext Book of Pulmonary Artery Catheterization, 92:798

Local and Regional Anesthesia; From the *Fundamentals of Anesthesia and Acute Medicine Series*, 92:1620Perioperative Medicine; Volume 18, Number 3 of *Anesthesiology Clinics of North America*, 92:561

Practical Echocardiography: The Heart of the Matter. Part 1, Basics, 92:285

Practical Management of Pain, 3rd Edition, 92:561

Proceedings of the Fourth International Symposium on the History of Anaesthesia, 92:1078

Regional Anesthesia; Volume 12, Number 2 (April 2000) of *Problems in Anesthesia*, 92:1078Regional Anesthesia; Volume 18, Number 2 of *Anesthesiology Clinics of North America*, 92:284

Textbook of Neuroanesthesia and Critical Care, 92:286

Transesophageal Echocardiography in Anesthesia, 92:562

Brachial plexus block, see Anesthetic techniques**Bradycardia**, see Heart**Brain**

- amino acids, extracellular, ischemia-induced accumulation, effects of 17 β -estradiol, 92:1520
- area postrema, effects of propofol, rats, 92:934
- auditory information processing, during adequate propofol anesthesia, 92:1210
- blood flow
 - hyperemic response to halothane, augmentation by endotoxin (letter), 92:1075
 - velocity, effect of CPAP, awake volunteers, 92:415
 - velocity, effects of sevoflurane vs halothane, children, 92:891
- cognitive function
 - decline after hypothermic CPB, and high jugular bulb venous oxygen saturation, 92:1370

- effects of small-dose ketamine on propofol sedation, 92:1465
- edema, outcome from head trauma plus hemorrhage, rats, 92:669
- epilepsy, epileptogenic property of sevoflurane (letter), 92:1359
- hemodynamic responses, desflurane vs sevoflurane (letter), 92:282
- hemorrhage, head trauma plus, outcome, rats, 92:669
- intracranial pressure, effects of 30% and 60% xenon inhalation, rabbits, 92:1245
- ischemia, protective effects of 17 β -estradiol, 92:1520
- memory, during adequate propofol anesthesia, 92:1210
- neoplasm, awake craniotomy, 92:89
- outcome from head trauma plus hemorrhage, rats, 92:669
- posterior cingulate and retrosplenial cortices, ketamine-induced c-Fos expression, effects of xenon and N₂O, rats, 92:362
- subarachnoid hemorrhage, microvascular endothelial dysfunction, rats, 92:990
- trauma
 - head-injured patients, hypertonic saline and antidiuretic hormone, 92:641
 - outcome from head trauma plus hemorrhage, rats, 92:669

Brain stem

- embolism, sudden respiratory arrest, patient undergoing endovascular abdominal aortic aneurysm repair, 92:335
- medial pontine reticular formation, supraspinal and spinal action of cholinomimetics, 92:1307

Brain surgery, see Surgery**Breast surgery**, see Surgery**Bronchoscopy**, see Surgery**Bupivacaine**, see Anesthetics, local**Calcium**

- extracellular, protamine and myocardial contractility, rats, 92:817
- intracellular
 - Ca²⁺ sensitivity, airway smooth muscle, effects of ethanol, 92:767
 - inhibition of Ca²⁺ transients by morphine, 92:602

Calcium channel blockers, see Pharmacology**cAMP**, see Heart, cyclic adenosine monophosphate**Carbachol**, see Pharmacology**Carbamazepine**, see Pharmacology

Carbon dioxide

absorbents

- carbon monoxide poisoning from hemolysis, 92:1609
- composition (letter), 92:1356
- end-tidal CO₂ monitoring during weaning from CPB, patients without significant lung disease, 92:306
- tracing configuration, dependence on sample size (letter), 92:1357
- vs transcutaneous CO₂ monitoring, during mechanical ventilation, older children, 92:306
- transcutaneous CO₂ monitoring, vs end-tidal CO₂ monitoring, during mechanical ventilation, older children, 92:306

Carbon monoxide, concentrations in breathing circuit, during low-flow anesthesia, effects of smoking status and body size, 92:542

Carbon monoxide poisoning, see Complications

Carboxyhemoglobin, see Blood

Carcinoma, see Liver

Cardiac arrest, see Complications; Heart

Cardiopulmonary bypass, see Surgery

Cardiopulmonary resuscitation (CPR)

- active compression-decompression (ACD), with inspiratory threshold valve, effects of PEEP, 92:967
- after incremental overdosage with lidocaine, bupivacaine, levobupivacaine, and ropivacaine, anesthetized dogs, 92:37
- endobronchial lidocaine administration, 92:1505
- experimental, effects of endogenous and exogenous vasopressin, 92:1499
- fluid, Pentalyte vs lactated Ringer's solution, effects on release of heparinoids and thrombotic mediators, rabbits, 92:314
- intraoperative, failed, survival after, case of Lazarus syndrome, 92:690
- neurological recovery, blood replacement with varying hematocrit levels, porcine model of profound hypothermic circulatory arrest, 92:329
- pulmonary blood flow and PETCO₂, 92:306

Cardiotoxicity, see Toxicity

Cardiovascular anesthesia, see Anesthesia

Carnitine palmitoyltransferase I

deficiency, see Complications

Catheter perforation, see Complications

Catheters, see Equipment

Caudal anesthesia, see Anesthetic techniques

Cell salvage bowls, see Equipment

Central nervous system

- effects of small-dose ketamine on propofol sedation, 92:1465
- inhibition of human $\alpha 7$ nicotinic acetylcholine receptor by thiopental, 92:930

Cerebrospinal fluid (CSF), leak, with caudal blood patch, 92:1169

Cervical epidural steroid injection, see Interventional pain management techniques

Cervical laminectomy, see Surgery

Cervical plexus block, see Anesthetic techniques

Cervical spine disposition, see Measurement techniques

C-Fos, see Protein

Channels

- ion
 - bupivacaine enantiomers on Ca²⁺ channels, 92:239
 - Ca²⁺ channel blockers and morphine, 92:239
 - ligand-gated, ATP and NMDA receptor interactions, 92:134
- potassium
 - blockage of one class, alteration of effectiveness of halothane in brain circuit, *Drosophila*, 92:535
 - mitochondrial ATP-sensitive, activation, protection of stunned myocardium, 92:1139
 - sodium, propofol block of muscle sodium channels, 92:1192

Chemotherapy, cisplatin, patient receiving, diffuse brachial plexopathy after interscalene blockade, pharmacological double crush syndrome, 92:249

Chest tomography, see Monitoring

Cholecystectomy, see Surgery

Cholesterol, apolipoprotein E, polymorphisms, and age at first CABG, 92:824

Chronic methylprednisolone infusion, see Anesthetic techniques

Chronic obstructive pulmonary disease, see Complications

Circle absorber system, see Equipment

Cisatracurium, see Neuromuscular relaxants

Cisplatin, see Chemotherapy

Cleft palate repair, see Surgery

Clonidine, see Pharmacology;

Premedication

Coagulation

antithrombin activity, low preoperative, effect on response to heparin, cardiac surgery patients, adults vs infants, 92:66

antithrombin III, MAX-ACT, patients undergoing extracorporeal circulation, 92:578

aprotinin, and pulmonary thromboembolism during orthotopic liver transplantation, 92:1418

epsilon-aminocaproic acid (ϵ -ACA), concentrations, in adults undergoing CPB, lack of effect of gender, 92:1384

heparin

-induced thrombocytopenia, patients with, use of lepirudin for anticoagulation during major vascular surgery, 92:344

MAX-ACT, patients undergoing extracorporeal circulation, 92:578

onset time, 92:1391

protamine and myocardial contractility, rats, 92:817

heparinoids, release, and Pentalyte, rabbits, 92:314

hypercoagulability, prolonged, after major abdominal surgery, thromboelastography for monitoring, 92:572

lepirudin, use for anticoagulation during major vascular surgery, patients with heparin-induced thrombocytopenia, 92:344

major orthopedic surgery, effects of Voluven vs HAES-steril, 92:855

platelet function

effect of magnesium, parturients with preeclampsia, 92:1257

effects of hydroxyethyl starches of varying molecular weights, 92:1402

protamine, effects on myocardium, modulation by extracellular calcium, rats, 92:817

protein function, effect of NO, thromboelastographic assessment, rabbits, 92:320

Sonoclot analysis, healthy children, 92:904

thromboelastography

effects of magnesium, 92:1257

for monitoring prolonged hypercoagulability after major abdominal surgery, 92:572

intravascular volume replacement and coagulation, 92:565

next step (editorial), 92:563

tranexamic acid levels during CPB, 92:1131

- Codeine**, see Analgesics
- Coffin-Siris syndrome**, see Complications
- Cognitive function**, see Brain
- Cold pressor test**, see Measurement techniques
- Colforsin daropate**, see Pharmacology
- Colloids**, see Fluid balance
- Colonoscopy**, see Surgery, colonic
- Coma**, see Complications
- Combined spinal-epidural analgesia**, see Anesthetic techniques
- Complex regional pain syndrome**, see Pain
- Complications**
- acute fatty liver of pregnancy, patient with, perioperative care, 92:1268
 - acute lung injury, patients with, breath termination criterion during pressure support ventilation, 92:161
 - acute respiratory distress syndrome, patients with, breath termination criterion during pressure support ventilation, 92:161
 - acute respiratory failure
 - gas exchange during CPAP and APRV, 92:950
 - ONO1714 and diaphragm in sepsis, 92:959
 - alveolar hemorrhage, negative pressure post-tracheal extubation, 92:273
 - arytenoid dislocation, while using McCoy laryngoscope, 92:1347
 - awareness
 - explicit intraoperative recall at BIS of 47, 92:808
 - search for reliable awareness monitor (editorial), 92:801
 - bilateral vocal cord palsy, after ventricular drainage, child, 92:358
 - bleeding, tranexamic acid levels during CPB, 92:1131
 - bradycardia, indigo carmine-induced, during general anesthesia, 92:276
 - breakage, epidural catheters, 92:246
 - carbon monoxide poisoning, from hemolysis, 92:1609
 - cardiac arrest, during spinal anesthesia, common mechanisms and strategies for prevention, 92:252
 - carnitine palmitoyltransferase I deficiency, child with, postoperative coma, 92:646
 - catheter perforation, venous double cannulation (letter), 92:558
 - chronic obstructive pulmonary disease, severe, patients with, thoracic epidural anesthesia and breathing, 92:1015
 - Coffin-Siris syndrome, patient with, difficult airway, 92:554
 - coma, postoperative, child with carnitine palmitoyltransferase deficiency, 92:646
 - compressive cervical hematoma, intravenous ketoprofen in thyroid and parathyroid surgery, 92:1052
 - congenital heart disease
 - infants and children, safety and efficacy of sevoflurane, 92:1152
 - infants with, volume ventilation, comparison of two ventilators, 92:76
 - constipation, oral ibuprofen vs codeine/acetaminophen, 92:1470
 - debris, elimination from partly filled cell salvage bowls, 92:1137
 - dysphagia, safety of intraoperative TEE, 92:1126
 - dystocia, intensity of labor pain and cesarean delivery, 92:1524
 - electrocautery and monitoring devices, 92:384
 - epidural catheter entrapment, 92:1532
 - esophageal injury, and unsuccessful TEE placement, 92:870
 - gastric insufflation, with ProSeal laryngeal mask, 92:1614
 - gastrointestinal trauma, safety of intraoperative TEE, 92:1126
 - hemophilia, patient with inhibitors to factor VIII, 92:785
 - human T lymphotropic virus type I-associated myelopathy, spinal anesthesia (letter), 92:1618
 - hypertension, tourniquet-induced, ketamine and, 92:1286
 - hyperventilation syndrome, ineffectiveness of propofol, 92:781
 - hypotension
 - angiotensin II antagonists and anesthetic induction, 92:26
 - severe, vaginal indigo-carmin induced (letter), 92:556
 - hypoxemic respiratory failure, life-threatening, after repair of aortic dissection, successful treatment, 92:872
 - intravascular injection, accidental, of lidocaine-bupivacaine mixture, and T-wave increase, 92:915
 - ischemia-reperfusion injury
 - lactate and indocyanine green elimination rate in cirrhosis, 92:1064
 - lung, protective effect of acadesine, 92:590
 - PentaLyte resuscitation, heparinoids, and thrombotic mediators, rabbits, 92:314
 - Klippel-Feil syndrome, patient with, thoracic epidural anesthesia for bilateral reduction mammoplasty, 92:514
 - lactic acidosis, olprinone and endotoxemia, 92:676
 - Larsen syndrome, parturient with, obstetric anesthetic management, 92:1266
 - Lazarus syndrome, survival after failed intraoperative resuscitation, 92:690
 - methemoglobinemia
 - after large-dose infusions of diaspirin cross-linked hemoglobin, 92:44
 - benzocaine-induced, factors influencing Hurricaine delivery, 92:379
 - myelopathy, human T lymphotropic virus type I-associated, spinal anesthesia (letter), 92:1618
 - neuropathic sensory disorders, chronic, effects of soy-containing diet, rats, 92:1029
 - obesity, morbid, knees-to-abdomen "Heimlich maneuver" (letter), 92:1619
 - Ortner's cardiovocal syndrome, cause of postoperative left vocal cord dysfunction, 92:1071
 - overinflation, air leaks, T_{1/2} of N₂O-filled cuffs deflation, 92:145
 - paradoxical fat embolism syndrome, fatal case, detection by intraoperative TEE, 92:688
 - paraplegia, reversible, associated with lumbar epidural analgesia and thoracic vertebral metastasis, 92:1316
 - Parkinson's disease, anesthesia for stereotactic frame placement, 92:424
 - paroxysmal nocturnal hemoglobinuria, patient with, continuous infusion of prostaglandin E₁ and epidural analgesia (letter), 92:1618
 - patent foramen ovale, detection using transnasal TEE, 92:1111
 - pelvic hematoma, after ilioinguinal nerve block for orchialgia, 92:1048
 - peripheral neuropathy, after interscalene blockade, patient receiving cisplatin therapy, pharmacological double crush syndrome, 92:249
 - pneumocephalus, after cervical epidural steroid injection, 92:1576
 - pneumonia, ventilator-associated, early diagnosis, evaluation of Gram stain in protected bronchoalveolar lavage fluid, 92:442

- polycythemia rubra vera, patient with, epidural anesthesia for cesarean section, 92:1535
- postdural puncture headache
- CSF leak with caudal blood patch, 92:1169
 - obstetric patients (letter), 92:1616
- preeclampsia, parturients with
- effect of magnesium on coagulation, 92:1257
- epidural anesthesia and polycythemia, 92:1535
- preoperative risk scores, efficacy and resource utilization of remifentanyl and fentanyl, fast-track CABG surgery, 92:1094
- recurrent laryngeal nerve palsy, and TEE probe, 92:1422
- reexpansion pulmonary edema, after thoracoscopic mediastinal tumor resection, 92:1416
- respiratory events, and failure of chain-link mechanism on anesthetic machine, 92:913
- shivering, postanesthetic, treatment, comparison between meperidine, clonidine, and urapidil, 92:257
- smoking, status, effects on carbon monoxide concentrations in breathing circuit during low-flow anesthesia, 92:542
- sore throat, pressure in E-T cuffs made of new material, 92:140
- spino-subcutaneous fistula, after bone marrow harvest, successful treatment with epidural blood patch, 92:1050
- superior laryngeal nerve injury, patient with, severe laryngospasm at tracheal extubation, 92:271
- systemic lupus erythematosus, myelopathy and laminectomy, 92:1254
- temporomandibular joint
- abnormal, cause of difficult tracheal intubation, 92:783
 - disorders, failure of local anesthesia to block mustard-oil-induced temporomandibular inflammation, 92:1035
- thrombocytopenia, heparin-induced, patients with, use of lepirudin for anticoagulation during major vascular surgery, 92:344
- thromboembolism, pulmonary
- antifibrinolytic therapy during orthotopic liver transplantation (letter), 92:559
 - aprotinin administration during orthotopic liver transplantation, 92:1418
 - transient neurologic symptoms after spinal anesthesia with lidocaine, obstetric patients, 92:405
 - lidocaine-induced, postcesarean section, 92:401
 - lidocaine neurotoxicity, obstetric patients (editorial), 92:287
 - vertebral metastasis, reversible paraplegia and epidural analgesia, 92:1316
 - wound infections, dexamethasone in ambulatory anesthesia, 92:85
- Compound A**, see Metabolism
- Compressive cervical hematoma**, see Complications
- Computed tomography**, see Imaging techniques
- Computer applications**
- anesthesia information management system, prediction of antiemetic rescue treatment, PACU, 92:1203
 - discrete-event computer simulation, OR utilization, 92:1215
 - Monte-Carlo simulation, statistical analysis of impact of delays in discharge from PACU on total patient care hours, 92:1222
- Congenital heart disease**, see Complications
- Conscious sedation**, see Anesthetic techniques
- Constipation**, see Complications
- Constrictive pericarditis**, see Heart
- Continuous infusion**, see Anesthetic techniques
- Continuous positive airway pressure**, see Ventilation
- Cooximetry**, see Measurement techniques
- Coronary artery disease**, see Complications
- Corrugator supercilii**, see Muscles
- Cost-benefit analysis**, see Economics
- Cough**, chemically induced, after bolus fentanyl injection, 92:1442
- CPAP**, see Ventilation, continuous positive airway pressure
- Craniotomy**, see Surgery
- Creatinine**, see Kidney
- Critical care and trauma**, see Anesthesia
- Crystalloids**, see Fluid balance
- Cytotoxicity**, see Toxicity
- Data management system**, risk factors, multivariate analysis, 92:1203
- Debris**, see Complications
- Desflurane**, see Anesthetics, volatile
- Dexamethasone**, see Anesthetics, intravenous; Pharmacology
- Dexmedetomidine**, see Anesthetics, intravenous
- Dextromethorphan**, see Pharmacology
- Diaphragm**, see Muscles; Ventilation
- Diaspirin cross-linked hemoglobin**, see Pharmacology
- Diet**, soy, suppression of chronic neuropathic sensory disorders, rats, 92:1029
- Diltiazem**, see Pharmacology
- Drug addiction**, patient recovering from, special issues for anesthesiologist, 92:1601
- Durshordwe, Clarence J.**, 92:1349
- Dysphagia**, see Complications
- Dyspnea**, see Ventilation
- Dystocia**, see Complications
- Ear**, auricular acupuncture, potential treatment for anxiety, 92:548
- Economics and health systems research**
- cost per dose of succinylcholine, 92:693
 - issues of concern for aging anesthesiologist, 92:1487
 - OR utilization, 92:1215
 - OR weekday staffing analysis, 92:1493
 - PACU discharge analysis, 92:1222
 - PACU staffing analysis, 92:947
 - patients' willingness to pay for antiemetic, 92:393
 - scheduling delays between cases, 92:943
 - systematic reviews in literature, 92:700
- ECT**, see Anesthesia, electroconvulsive therapy
- Edrophonium**, see Antagonists, neuromuscular relaxants
- Education**, simple algorithm for tracheal intubation, 92:517
- Elderly**, see Age factors
- Electricity**, radiofrequency transmission, to monitoring devices in OR, simulation study, 92:384
- Electroanalgesia**, see Anesthetic techniques
- Electroconvulsive therapy**, see Anesthesia
- Electroencephalography**, see Monitoring
- Electromyography**, see Measurement techniques; Monitoring
- Embolism**
- air, transnasal TEE and foramen ovale patency, 92:1111
 - brain stem, sudden respiratory arrest, patient undergoing endovascular abdominal aortic aneurysm repair, 92:335
 - fat embolic events during hip arthroplasty, 92:49

Embolism—continued

- fat elimination from autologous blood, 92:341
- paradoxical fat embolism syndrome, fatal case, detection by intraoperative TEE, 92:688
- pulmonary, and aprotinin administration, during orthotopic liver transplantation, 92:1418

Emergence, from anesthesia

- direct current auditory evoked potentials, 92:154
- intranasal fentanyl, 92:154

Endobronchial lidocaine, see Anesthetic techniques**Endothelium, microvascular endothelial dysfunction, after subarachnoid hemorrhage, rats, 92:990****Endotoxemia, see Infection****Endotracheal tubes, see Equipment****End-tidal CO₂, see Monitoring****Enflurane, see Anesthetics, volatile****Enzymes**

- carnitine palmitoyltransferase, deficiency, child with, postoperative coma, 92:646
- phosphodiesterase type III, inhibition before aortic cross-clamping preserves intramyocardial adenosine monophosphate during CPB, 92:1377

Epidural, see Anesthetic techniques**Epidural blood patch, see Anesthetic techniques****Epilepsy, see Brain****Epinephrine, see Pharmacology****Epsilon-aminocaproic acid, see Coagulation****Equipment**

- anesthetic machines, Ohmeda Excel 210 SE, failure of chain-link mechanism, 92:913
- arterial lines, securing (letter), 92:1355
- blood filters, fat elimination from autologous blood, 92:341
- catheters
 - epidural, "blocked," another cause (letter), 92:1617
 - epidural, breakage, arrow reinforced catheter vs other nonreinforced catheters, 92:246
 - epidural, periosteal entrapment in intrathecal space, 92:1532
 - epidural, tunneled, for prolonged analgesia, pediatric patients, 92:1432
 - interscalene brachial plexus, difficult insertion (letter), 92:1074
 - intravenous, new, and transtracheal jet ventilation (letter), 92:1074

- perforation, complication of venous double cannulation (letter), 92:558
- pulmonary artery, and tricuspid or pulmonic valvular regurgitation, 92:1117

cell salvage bowls, partly filled, debris elimination, 92:1137**circle absorber system, preoxygenation, tidal volume vs deep breaths, 92:1337****endotracheal tubes**

- cuff, N₂O-filled, deflationary phenomenon after cessation of N₂O, 92:145

exchange, complexity of procedure (letter), 92:796**Profile Soft-Seal Cuff, N₂O gas-barrier properties, 92:140****vs LMA, as route for endobronchial lidocaine administration, 92:1505****esophageal detector devices, efficacy in verifying tracheal tube placement, 92:375****fiberoptic bronchoscope, -aided tracheal intubation, two-person technique, patient with long and narrow retropharyngeal air space, 92:1611****glutaraldehyde fume hood, creation of individual soaking compartments (letter), 92:1073****intravenous lines, securing (letter), 92:1355****intubating laryngeal mask airway (ILMA), vs awake fiberoptic intubation, patients with difficult airways, 92:1342****laryngeal mask airway (LMA)****extubation, MAC of enflurane, deeply anesthetized children, 92:72****ProSeal, gastric insufflation with, 92:1614****vs endotracheal tube, as route for endobronchial lidocaine administration, 92:1505****laryngoscope**

- blades, articulating, vs external laryngeal manipulation, videographic analysis of laryngeal exposure, 92:267

blades, individual soaking compartments within glutaraldehyde fume hood (letter), 92:1073**laryngoscopy and spinal cord space deformation, 92:1331****McCoy laryngoscope, arytenoid dislocation, 92:1347****masks, use of, during neuraxial block placement (letters), 92:279****microneedles, microfabricated, lack of pain associated with, 92:502****negative pressure warming, efficacy, 92:261****nonmetered sprays, factors influencing Hurricaine delivery, 92:379****pressure bags, for securing intravenous and arterial lines (letter), 92:1355****pulse oximeter, wave form, ear and finger, different responses to cold pressor test, 92:1483****tourniquet, -induced arterial pressure increase, prevention with preoperative small-dose ketamine, 92:1286****unlighted stylet, tracheal intubation (letter), 92:1355****ventilators**

- associated pneumonia, early diagnosis, evaluation of Gram stain in protected bronchoalveolar lavage fluid, 92:442

comparison, volume ventilation of infants with congenital heart disease, 92:76**ventricular-peritoneal shunt, insertion, followed by bilateral vocal cord palsy, child, 92:358****Esmolol, see Pharmacology****Esophageal detector devices, see Equipment****Esophageal injury, see Complications****Esophagectomy, see Surgery****17 β -Estradiol, see Hormones****Ethanol, see Pharmacology****Extended femoral nerve sheath block, see Anesthetic techniques****Extracorporeal shock wave lithotripsy, see Surgery****Extubation**

- immediate or early, after coronary artery bypass surgery (letter), 92:1073

tracheal

- negative pressure post-tracheal extubation alveolar hemorrhage, 92:273

severe laryngospasm, patient with superior laryngeal nerve injury, 92:271**Eye, see Surgery, ophthalmic****Fascia iliaca compartment block, see Anesthetic techniques****Fast-tracking, see Anesthetic techniques****Fat, see Embolism****Femoral nerve block, see Anesthetic techniques****Fiberoptic bronchoscope, see Equipment****FK506, see Pharmacology, tacrolimus****Fluid balance**

albumin, treatment with, effect on
neurologic status and brain edema,
head trauma plus hemorrhage, rats,
92:669

colloids, vs crystalloids, intravascular
volume before obstetric spinal
anesthesia, 92:997

crystalloids, vs colloids, intravascular
volume before obstetric spinal
anesthesia, 92:997

gelatin
intravascular volume replacement and
coagulation, 92:565
thromboelastography, next step
(editorial), 92:563

hetastarch
intravascular volume before obstetric
spinal anesthesia, 92:997
treatment with, effect on neurologic
status and brain edema, head
trauma plus hemorrhage, rats,
92:669

hydroxyethyl starch
hemostatic changes, influence of ABO
blood group, 92:1396
intravascular volume replacement and
coagulation, 92:565
novel, Voluven, vs HAES-steril' effects
on coagulation in major orthopedic
surgery, 92:855
thromboelastography, next step
(editorial), 92:563
varying molecular weights, effects on
platelet function, 92:1402

hypertonic saline
treatment with, effect on neurologic
status and brain edema, head
trauma plus hemorrhage, rats,
92:669
vs lactated Ringer's solution, and
antidiuretic hormone in severe head
injury, 92:641

lactated Ringer's solution
infusion, effects on cardiac output
changes after spinal anesthesia,
92:710
intravascular volume before obstetric
spinal anesthesia, 92:997
vs hypertonic saline, and antidiuretic
hormone in severe head injury,
92:641
vs PentaLyte, effects on release of
heparinoids and thrombotic
mediators, rabbits, 92:314

PentaLyte, vs lactated Ringer's solution,
effects on release of heparinoids
and thrombotic mediators, rabbits,
92:314

Foramen ovale, see Heart

GABA_A receptors, see Receptors

Gases, nonanesthetic, nitric oxide (NO)
diaspirin cross-linked hemoglobin and
vascular tone, 92:324
donors, DETANONOate, effect on
coagulation protein function,
thromboelastographic assessment,
rabbits, 92:320
inhaled, effects on oxygenation during
one-lung ventilation, 92:842
production, from activated murine
macrophage cell line, effects of
lidocaine, 92:128
selective iNOS inhibition, sepsis, rats,
92:681
synthase
induction by endotoxin, and cerebral
hyperemic response to halothane
(letter), 92:1075
inhibitors, inducible, ONO1714,
attenuation of sepsis-induced
diaphragmatic dysfunction,
hamsters, 92:959
vasomotion in subarachnoid
hemorrhage, rats, 92:990

Gastric insufflation, see Complications

Gastrointestinal tract, jejunal mucosal
perfusion, during mild hypothermic
CPB, 92:5

Gastrointestinal trauma, see
Complications

Gelatin, see Fluid balance

Gender, and ϵ -ACA concentrations, adults
undergoing CPB, 92:1384

General anesthesia, see Anesthetic
techniques

Gene therapy, with vascular endothelial
growth factor, for inoperable
coronary artery disease, anesthetic
management and results, 92:19

Genotoxicity, see Toxicity

Glucocorticoids, see Pharmacology

Glutamate receptors, see Receptors

Glutaraldehyde fume hood, see
Equipment

Glycine receptors, see Receptors

Gynecologic surgery, see Surgery

Hand surgery, see Surgery

Heart
aortic valve
area, hemodynamic changes,
implications for Doppler cardiac
output determinations, 92:584
prosthesis, acute thrombosis,
diagnosis by TEE, 92:584
arrest
effects of PEEP during active
compression-decompression CPR

with inspiratory threshold valve,
92:967
efficacy of esophageal detector devices
in verifying tracheal tube
placement, 92:375
endogenous vs exogenous vasopressin
during experimental CPR, 92:1499
arrhythmias, post-CABG, effect of insulin,
92:810
atrial fibrillation, post-CABG, effect of
insulin, 92:810
bradycardia, indigo carmine-induced,
during general anesthesia, 92:276
cardiac output
after spinal anesthesia, 92:710
determinants of propofol induction of
anesthesia dose, 92:656
hemodynamic changes in aortic valve
area, 92:584
prediction of postoperative
hypovolemia, 92:1146
stroke volume variation, fluids, and
craniotomy, 92:984
cardiovascular function, effects of prone
positioning, patients with acute
lung injury, 92:1226

catecholamines
fast-tracking after CABG (editorial),
92:1081
remifentanyl vs fentanyl in CABG,
92:1084

constrictive pericarditis, intraoperative
hemodynamic and
echocardiographic evaluation of
cardiac filling dynamics, 92:1424

contractility
effects of protamine, modulation by
extracellular calcium, rats, 92:817
ethanol and stunned myocardium,
dogs, 92:299
inhibition of Ca^{2+} transients by
morphine, 92:602

cyclic adenosine monophosphate
(cAMP), during CPB, and inhibition
of phosphodiesterase type III before
aortic cross-clamping, 92:1377

diastolic function, dysfunction as
predictive of difficult weaning from
CPB, 92:291

differential perfusion, during CPB, TEE
imaging of new aortic cannula,
92:338

enzyme concentration
fast-tracking after CABG (editorial),
92:1081
remifentanyl vs fentanyl in CABG,
92:1084

failure, after CABG, predicting risk of
death, 92:596

Heart—continued

- foramen ovale, patent, detection using transnasal TEE, 92:1111
- ischemia
 - detecting, evaluation of new two-electrode myocardial electrical impedance monitor, 92:12
 - ethanol and stunned myocardium, dogs, 92:299
 - gene therapy for inoperable coronary disease, 92:19
 - preischemic conditioning, preservation of intramyocardial cAMP during CPB, 92:1377
- myocardium, sevoflurane and mitochondrial K_{ATP} channels, 92:1139
- rate, during ECT, effects of diltiazem, 92:1327
- reflex tachycardia, absence after spinal anesthesia, elderly patients (letter), 92:1357

Heat shock protein 70, see Protein

Hematocrit, see Blood

Hemoglobin, see Blood

Hemoglobin-based oxygen carrier, see Blood

Hemolysis, see Blood

Hemophilia, see Complications

Hemorrhage

- alveolar, negative pressure post-tracheal extubation, 92:273
- diaspirin cross-linked hemoglobin, 92:44
- outcome from head trauma plus hemorrhage, rats, 92:669

Heparin, see Coagulation; Pharmacology

Heparinoids, see Coagulation

Hepatectomy, see Surgery

History, anesthesia, Clarence D. Durshordwe, 92:1349

Hoarseness, see Airway

Hormones

- antidiuretic, release, response to hypovolemia and/or sodium administration, children with severe head injury, 92:641
- 17 β -estradiol, protection of neurons from ischemic damage and attenuation of accumulation of extracellular excitatory amino acids, 92:1520

Hospital, preadmission, trauma, tramadol for pain relief, 92:1543

Human leukocyte antigen-DR (HLA-DR), see Monocytes

Human T lymphotropic virus type I-associated myelopathy, see Complications

Hyaluronidase, see Pharmacology

Hydrocarbons, see Mechanisms of anesthetic action

Hydroxyethyl starch, see Fluid balance

Hyperalgesia, see Pain

Hypercarbic ventilatory response, see Ventilation

Hypertension, see Blood pressure; Complications

Hyperventilation syndrome, see Complications

Hypotension, see Blood pressure; Complications

Hypothermia, see Temperature

Hypoxemic respiratory failure, see Complications

Hysterectomy, see Surgery

IARS, see International Anesthesia Research Society

Ibuprofen, see Analgesics

Ilioinguinal nerve block, see Anesthetic techniques

ILMA, see Equipment, intubating laryngeal mask airway

Imaging techniques

- computed tomography (CT), three-dimensional, and prediction of unanticipated difficult tracheal intubation (letter), 92:281
- ultrasonography, examination of axillary part of brachial plexus, 92:1271
- video imaging, assessment of neuromuscular blockade at larynx, 92:149

Immune system

- lymphocyte subsets, effects of epidural block, patients with and without pain, 92:463
- natural-killer cell activity, effects of epidural block, patients with and without pain, 92:463
- suppression
 - early on during surgery, 92:1322
 - FK506 and spinal cord ischemia, rats, 92:1237

Indigo carmine, see Pharmacology

Indocyanine green, see Measurement techniques

Infection

- endotoxemia, porcine model, effects of olprinone on hepatic vascular bed, 92:676
- nosocomial, ventilator-associated pneumonia, early diagnosis, evaluation of Gram stain in protected bronchoalveolar lavage fluid, 92:442
- sepsis

-induced diaphragmatic dysfunction, attenuation by ONO1714, hamsters, 92:959

neutrophil hydrogen peroxide production and adhesion molecule expression, effects of propofol vs midazolam, rats, 92:437

selective iNOS inhibition, prevention of hypotension and preservation of endothelium-dependent vasodilation, rats, 92:681

Inflammation

- adhesion molecules
 - and TGF- β 1, in ventilator-induced lung injury, 92:428
- expression, effects of propofol vs midazolam, abdominal sepsis, rats, 92:437
- macrophages, activation, and ventilator-induced lung injury, rats, 92:428
- neutrophils
 - hydrogen peroxide production, effects of propofol vs midazolam, abdominal sepsis, rats, 92:437
- infiltration, and ventilator-induced lung injury, rats, 92:428
- lysophosphatidic acid-induced priming agents, attenuation by local anesthetics, 92:1041
- synovial, plasma extravasation, inhibition by preemptive administration of antiinflammatory irrigation solution, knee, rats, 92:1301
- temporomandibular, mustard-oil-induced, failure of local anesthesia to block, 92:1035

Information systems, PACU, prediction of antiemetic rescue treatment, 92:1203

Insulin, see Pharmacology

Intensive care unit (ICU), propofol vs midazolam, mechanically ventilated critically ill patients, 92:975

Interactions, drug

- clonidine and lidocaine, 92:733
- fentanyl and levobupivacaine, 92:410
- ketorolac and sevoflurane, 92:1058
- levobupivacaine and fentanyl, 92:410
- lidocaine and clonidine, 92:733
- neuromuscular relaxants, 92:1199
- ondansetron and tramadol, 92:1319
- sevoflurane and ketorolac, 92:1058
- tramadol and ondansetron, 92:1319

International Anesthesia Research Society (IARS), history, 92:1349

International Society for Anaesthetic Pharmacology (ISAP), name change from SIVA (editorial), 92:1

- Internet**, technology and innovation,
Anesthesia & Analgesia online
(editorial), 92:1363
- Interscalene block**, see Anesthetic
techniques
- Interventional pain management**
techniques, cervical epidural steroid
injection, pneumocephalus after,
92:1576
- Intravascular injection**, see Complications
- Intravenous lines**, see Equipment
- Intravenous regional anesthesia**, see
Anesthetic techniques
- Intubation**
endotracheal, with thiopental/
succinylcholine or sevoflurane-N₂O,
comparative study, adults, 92:523
laryngeal, conventional vs balloon
laryngoscopy-guided, disposition of
cervical spine and deformation of
available cord space, 92:1331
tracheal
articulating blade vs laryngeal
manipulation, 92:267
awake fiberoptic intubation vs ILMA,
patients with difficult airways,
92:1342
difficult, as result of unsuspected
abnormality of temporomandibular
joint, 92:783
difficult, two-person technique for
fiberscope-aided intubation, 92:1611
difficult, unanticipated, prediction, use
of three-dimensional CT imaging
(letter), 92:281
difficult, use of McCoy laryngoscope,
and arytenoid dislocation, 92:1347
fiberscope-aided, two-person
technique, patient with long and
narrow retropharyngeal air space,
92:1611
simple algorithm, validation, 92:517
unlighted stylet (letter), 92:1355
verifying tracheal tube placement,
efficacy of esophageal detector
devices, 92:375
- Ion channels**, see Channels
- Ions**, magnesium
effect on coagulation and platelet
function, parturients with
preeclampsia, 92:1257
modulation of NMDA receptor function,
92:1173
modulation of NMDA receptor function,
interactions with volatile
anesthetics, 92:1182
- Ischemia**, see Aorta; Brain; Heart; Lung;
Spinal cord
- Ischemia-reperfusion injury**, see
Complications
- Jejunal mucosal perfusion**, see
Gastrointestinal tract
- Jugular venous oxygen saturation**, see
Monitoring
- Ketamine**, see Anesthetics, intravenous;
Pharmacology
- Kidney**
creatinine, allosteric modification of
oxygen delivery by hemoglobin,
92:615
damage, safety of ketorolac and
sevoflurane anesthesia, 92:1058
failure, patients with, effects of low-flow
sevoflurane and isoflurane
anesthesia, 92:650
function, effects of prone positioning,
patients with acute lung injury,
92:1226
- Klippel-Feil syndrome**, see Complications
- Knee surgery**, see Surgery
- Labor**, see Pain
- Lactated Ringer's solution**, see Fluid
balance
- Lactate levels**, see Measurement
techniques
- Lactic acidosis**, see Complications
- Laparoscopic surgery**, see Surgery
- Laparotomy**, see Surgery
- Larsen syndrome**, see Complications
- Laryngeal mask airway**, see Equipment
- Laryngoscope**, see Equipment
- Larynx**
laryngeal exposure, videographic
analysis, articulating laryngoscope
vs external laryngeal manipulation,
92:267
neuromuscular blockade, assessment,
video imaging, 92:149
- Laser Doppler flowmetry**, see Monitoring
- Lasers**, function, anesthetic interference
during excimer laser procedures,
children, 92:1444
- LASIK procedures**, see Surgery
- Lazarus syndrome**, see Complications
- Lepirudin**, see Coagulation
- Levobupivacaine**, see Anesthetics, local
- Lidocaine**, see Anesthetics, local
- Lipopolysaccharide**, hyporesponsiveness,
perioperative, immunosuppression
early on during surgery, 92:1322
- Lithotomy position**, see Position
- Liver**
acute fatty liver of pregnancy, patient
with, perioperative care, 92:1268
blood flow, effects of olprinone,
endotoxemia, porcine model, 92:676
carcinoma, lactate and indocyanine
green elimination rate in cirrhosis,
92:1064
failure, after acetaminophen and
sevoflurane exposure, child, 92:1446
- Liver transplantation**, see Surgery
- LMA**, see Equipment, laryngeal mask
airway
- Low-flow anesthesia**, see Anesthetic
techniques
- Lung**
blood flow, and PETCO₂, 92:306
hypertension, pulmonary arterial
pressure and TEE, 92:1364
injury
acute, patients with, effects of prone
positioning, 92:1226
ischemia-reperfusion, protective effect
of acadesine, 92:590
ventilator-induced, adhesion
molecules and TGF- β 1, 92:428
ischemia, -reperfusion injury, protective
effect of acadesine, 92:590
pulmonary edema
negative pressure post-tracheal
extubation alveolar hemorrhage,
92:273
reexpansion, after thoracoscopic
mediastinal tumor resection, 92:1416
pulmonary shunt, embolic events during
hip arthroplasty, 92:49
pulmonary thromboembolism and
antifibrinolytic therapy during
orthotopic liver transplantation
(letter), 92:559
- Lymphocyte subsets**, see Immune system
- MAC**, see Potency, minimum alveolar
concentration
- Macrophages**, see Inflammation
- Magnesium**, see Ions
- Malignant hyperthermia**, see Temperature
- Mammoplasty**, see Surgery
- Masks**, see Equipment
- mCD14**, see Monocytes
- McMechan, Francis Hoeffler**, founder of
IARS, 92:1349
- Measurement techniques**
activated clotting time
clinical onset of heparin, 92:1391
MAX-ACT, patients undergoing
extracorporeal circulation, 92:578
blood gas analysis, interference by
hemoglobin-based blood substitutes,
92:863
cardiac output, prediction of
postoperative hypovolemia, 92:1146

Measurement techniques—continued

- cervical spine disposition, conventional vs balloon laryngoscopy-guided laryngeal intubation, 92:1331
- circulating blood volume, use of hemoglobin-based oxygen carrier, rabbits, 92:609
- cold pressor test, different responses of ear and finger pulse oximeter wave form, 92:1483
- cooximetry, interference by hemoglobin-based blood substitutes, 92:863
- electromyography, ropivacaine vs bupivacaine for PCEA, 92:1024
- extracellular fluid, prediction of postoperative hypovolemia, 92:1146
- health status after anesthesia and surgery, questionnaire (letter), 92:281
- indocyanine green, elimination rate, and lactate, cirrhotic patients, 92:1064
- initial distribution volume of glucose, prediction of postoperative hypovolemia, 92:1146
- lactate levels, indocyanine green elimination rate, cirrhotic patients, 92:1064
- messenger RNA, lidocaine and iNOS in macrophage, 92:128
- micronucleus assay, genotoxicity of waste anesthetic gases, 92:118
- Northern blot analysis, lidocaine and iNOS in macrophage, 92:128
- oximetry, interference by hemoglobin-based blood substitutes, 92:863
- pain scores, Children's and Infants Postoperative Pain Score, perioperative analgesia and acetaminophen, 92:907
- percentage of glottic opening score, articulating blade vs laryngeal manipulation, 92:267
- proton-transfer-reaction mass spectrometry, online monitoring of air quality in PACU, 92:389
- sonography, study of diaphragm movement before and after cholecystectomy, 92:755
- thromboelastography
 - assessment of NO effect on coagulation protein function, rabbits, 92:320
 - intravascular volume replacement and coagulation, 92:565
 - next step (editorial), 92:563
 - Pentalyte resuscitation, heparinoids, and thrombotic mediators, rabbits, 92:314

- transcranial Doppler ultrasonography, effect of CPAP on cerebral blood flow velocity, awake volunteers, 92:415

- Western blot analysis, lidocaine and iNOS in macrophage, 92:128

Mechanisms of anesthetic action

- GABA_A block reverses isoflurane effects, 92:1578
- GABA_A glycine and MAC, 92:123
- GABA_A receptors and isoflurane MAC, 92:1585
- hydrocarbons, molecular determinants of anesthetic potency, 92:1477
- Shaker and halothane in fruit flies, 92:535

Memory, see Brain; Recall**Meperidine, see Analgesics; Pharmacology****Messenger RNA, see Measurement techniques****Meta-analysis, see Statistics****Metabolism, Compound A, sevoflurane in patients with renal failure, 92:650****Metamizol, see Analgesics****Methemoglobin, see Blood****Methemoglobinemia, see Complications****Methohexital, see Anesthetics, intravenous****Methylprednisolone, see Pharmacology, glucocorticoids****Microneedles, see Equipment****Micronucleus assay, see Measurement techniques****Midazolam, see Pharmacokinetics; Pharmacology; Premedication****Milrinone, see Pharmacology****Minimum alveolar concentration, see Potency, anesthetic****Mivacurium, see Neuromuscular relaxants****Monitored anesthesia care, see Anesthetic techniques****Monitoring**

- auditory evoked potentials, direct current, during wakefulness, anesthesia, and emergence from anesthesia, 92:154
- chest tomography, negative pressure post-tracheal extubation alveolar hemorrhage, 92:273
- devices, in OR, radiofrequency transmission, simulation study, 92:384
- electroencephalography
 - bispectral index, adequate propofol anesthesia and memory, 92:1210
 - bispectral index, as awareness monitor (editorial), 92:801
 - bispectral index, effect on anesthetic use and recovery, children anesthetized with sevoflurane in N₂O, 92:877

bispectral index, explicit

- intraoperative recall at BIS of 47, 92:808
- bispectral index, in pediatric anesthesia (editorial), 92:805
- effects of propofol sedation on sleep, 92:1232
- epileptiform EEG activity, elicited by sevoflurane, patients without epilepsy (letter), 92:1359
- electromyography, intramuscular vs surface, for determining diaphragmatic neuromuscular blockade, 92:106
- end-tidal CO₂
 - during weaning from CPB, patients without significant lung disease, 92:306
 - tracing configuration, dependence on sample size (letter), 92:1357
 - vs transcutaneous CO₂ monitoring, during mechanical ventilation, older children, 92:306
- intraoperative, in neuroanesthesia, national comparison between two surveys in Germany, 1991 and 1997, 92:166
- jugular venous oxygen saturation during and after CPB, 92:837
 - high values, association with cognitive decline after hypothermic CPB, 92:1370
- laser Doppler flowmetry, jejunal mucosal perfusion during mild hypothermic CPB, 92:5
- myocardial electrical impedance, two-electrode, for detecting myocardial ischemia, evaluation, 92:12
- nerve stimulation, brachial plexus block at humeral canal, 92:194
- peripheral venous pressure, neurosurgical patients, 92:172
- plethysmography, cold immersion, finger and ear, 92:1483
- pulmonary arterial pressure, intraoperative, estimation using pulsed Doppler TEE, 92:1364
- recurrent laryngeal nerve, during single-lung ventilation in esophagectomy, 92:662
- somatosensory evoked potentials (SSEPs)
 - intraoperative, remifentanyl- and fentanyl-based anesthesia, 92:1510
- spinal, inhibition by direct tramadol application on sciatic nerve, rats, 92:1547

- stroke volume variation, as predictor of
fluid responsiveness, patients
undergoing brain surgery, 92:984
- transcutaneous CO₂, vs end-tidal CO₂
monitoring, during mechanical
ventilation, older children, 92:306
- transesophageal echocardiography (TEE)
detection of embolic events during
cemented total hip arthroplasty,
92:49
- diagnosis of thrombosis of aortic
valve prosthesis, 92:1123
- diastolic dysfunction as predictive of
difficult weaning from CPB, 92:291
- evaluation of cardiac filling dynamics,
constrictive pericarditis, 92:1424
- hemodynamic changes in aortic valve
area, 92:584
- imaging of new aortic cannula for
differential perfusion during CPB,
92:338
- infants with compromised ventilation,
proper probe positioning (letter),
92:1076
- intraoperative, basic, performed by
experienced anesthesiologists,
adequacy, 92:1103
- intraoperative, detection of
paradoxical fat embolism syndrome,
92:688
- intraoperative, safety, 92:1126
- probe, failed placement, and recurrent
laryngeal nerve palsy, 92:1422
- pulsed Doppler, estimated of
pulmonary arterial pressure, 92:1364
- transnasal, detection of interatrial flow
patency, awake and anesthetized
patients, 92:1111
- unsuccessful placement, and
esophageal pathology, 92:870
- Monocytes**, mCD14 and HLA-DR
expression, and
immunosuppression early on
during surgery, 92:1322
- Monte-Carlo simulation**, see Statistics
- Mood**, postoperative, effects of small-dose
ketamine on propofol sedation,
92:1465
- Morphine**, see Analgesics
- Muscles**
corrugator supercilii, intramuscular vs
surface electromyography, for
determining neuromuscular
blockade, 92:106
- diaphragm
intramuscular vs surface
electromyography, for determining
neuromuscular blockade, 92:106
- kinetics, movement before and after
cholecystectomy, sonographic study,
92:755
- orbicularis oculi, intramuscular vs
surface electromyography, for
determining neuromuscular
blockade, 92:106
- skeletal, propofol block of muscle
sodium channels, 92:1192
- smooth
airway, dose-response study of
anticholinesterase drugs, rats, 92:100
- airway, effects of ethanol on Ca²⁺
sensitivity, 92:767
- uterus, contractility, effects of opioids,
92:1006
- Music**, sensory stimuli and anxiety,
children undergoing surgery, 92:897
- Myelopathy**, see Complications
- Myocardial electrical impedance**, see
Monitoring
- Natural-killer cell activity**, see Immune
system
- Nefopam**, see Analgesics
- Neostigmine**, see Antagonists,
neuromuscular relaxants;
Pharmacology
- Nephrotoxicity**, see Toxicity
- Nerve**, sciatic
anterior sciatic nerve block and leg
rotation, 92:460
- direct tramadol application, inhibition of
spinal SSEPs, rats, 92:1547
- division, in popliteal fossa, anatomical
implications for popliteal nerve
blockade, 92:215
- Nerve blocks**, see Anesthetic techniques
- Nerve injury**, chronic pain suppression
with dietary soy, 92:1029
- Nerve stimulation**, see Monitoring
- Neuraxial block**, see Anesthetic techniques
- Neuromuscular relaxants**
cisatracurium, severe anaphylactic
reaction, child, 92:648
- mivacurium, diaphragmatic
neuromuscular blockade,
monitoring, intramuscular vs
surface electromyography, 92:106
- rapacuronium
as alternative to succinylcholine for
ECT, 92:1171
- intubating dose, duration of action of
subsequent maintenance dose of
vecuronium, 92:1199
- succinylcholine
cost identification analysis, 92:693
- intubating dose, duration of action of
subsequent maintenance dose of
vecuronium, 92:1199
- neuromuscular blockade at larynx,
assessment, video imaging, 92:149
- rapacuronium as alternative to, for
ECT, 92:1171
- thiopental/succinylcholine, vs
sevoflurane-N₂O, endotracheal
intubation, adults, 92:523
- vecuronium
duration of action after intubating
dose of rapacuronium, vecuronium,
or succinylcholine, 92:1199
- intubating dose, duration of action of
subsequent maintenance dose of
vecuronium, 92:1199
- Neuropathic pain**, see Pain
- Neuropathic sensory disorders**, see
Complications
- Neurosurgical anesthesia**, see Anesthesia
- Neurotoxicity**, see Toxicity
- Neutrophils**, see Inflammation
- Nicotinic receptors**, see Receptors
- Nitric oxide**, see Gases, nonanesthetic
- Nitrous oxide**, see Anesthetics, gases
- NMDA receptors**, see Receptors
- Nociception**, see Pain
- Nonmetered sprays**, see Equipment
- Northern blot analysis**, see Measurement
techniques
- Nutrition**, early oral nutrition, multimodal
postoperative rehabilitation
program, 92:1594
- Obesity**, see Complications
- O blood group**, see Blood groups
- Obstetric anesthesia**, see Anesthesia
- Occupational hazards**
air quality, PACU, online monitoring,
92:389
- issues of concern for aging
anesthesiologists, 92:1487
- Olprinone**, see Pharmacology,
phosphodiesterase III inhibitors
- Ondansetron**, see Pharmacology
- One-lung ventilation**, see Ventilation
- ONO-8711**, see Pharmacology
- Operating rooms (OR)**
delays
PACU discharge analysis, 92:1222
- PACU staffing analysis, 92:947
- scheduling between cases, using
upper prediction bounds for case
durations, 92:943
- monitoring devices, radiofrequency
transmission, simulation study,
92:384
- occupational exposure to inhaled
anesthetics, high-level vs low-level,
association with genotoxicity,
micronucleus assay, 92:118

Operating rooms (OR)—continued

utilization, high, surgical suites, impact on revenue of increasing patient volume, 92:1215

weekday staffing analysis, 92:1493

Ophthalmic surgery, see Surgery**Opioid receptors, see Receptors****Opioids**

effects on isolated human pregnant uterine muscles, 92:1006

requirement, after cesarean delivery, effect of intravenous ketorolac, 92:1010

Orbicularis oculi, see Muscles**Orbital block, see Anesthetic techniques****Orchialgia, see Pain****Ortner's cardiopulmonary syndrome, see Complications****Outcomes**

predicted risk of fatal heart failure, 92:596

simple algorithm for tracheal intubation, 92:517

survival after failed intraoperative resuscitation, 92:690

Oximetry, see Measurement techniques**Oxycodone, see Pharmacology****Oxygen**

inspired fraction of oxygen, increasing levels, effects on oxygenation during one-lung ventilation, 92:842

measurement, cooximetry interference by hemoglobin-based blood substitutes, 92:863

supplemental intraoperative, vs ondansetron, for prevention of PONV, 92:112

transport, allosteric modification of oxygen delivery by hemoglobin, 92:615

Oxygenation

during one-lung ventilation, thoracic surgery

effects of inhaled NO and increasing levels of inspired fraction of oxygen, 92:842

thoracic epidural anesthesia combined with general anesthesia, 92:848

preoxygenation, tidal volume vs deep breaths, 92:1337

Oxyhemoglobin, see Blood**PACU, see Postanesthesia care unit****Pain**

acute, management, role of electroanalgesia, 92:505

allodynia, after partial sciatic nerve ligation, and soy-containing diet, rats, 92:1029

antinociception

interaction of clonidine and lidocaine, 92:733

supraspinal and spinal action of cholinomimetics, 92:1307

chronic

management, role of electroanalgesia, 92:505

suppression by dietary soy, rats, 92:1029

complex regional pain syndrome

glucocorticoid inhibition of hyperalgesia, 92:476

response to carbamazepine and sustained-release morphine, patients pretreated with spinal cord stimulation, 92:488

formalin-induced, analgesic interaction between intrathecal clonidine and glutamate receptor antagonists, rats, 92:725

heat pain threshold, effect of remifentanyl, volunteers, 92:369

hyperalgesia

acute tolerance to alfentanil, 92:1296

after partial sciatic nerve ligation, and soy-containing diet, rats, 92:1029

EP₁ antagonist and postoperative pain, 92:233

neuropathic, glucocorticoid inhibition, 92:476

thermal, effect of α 1- and α 2-adrenergic antagonists, 92:1558

inflammatory, and acute tolerance to alfentanil, 92:1296

labor

addition of morphine to fentanyl-bupivacaine spinal analgesia, 92:665

bupivacaine vs ropivacaine, potencies and probabilities (letter), 92:278

cesarean delivery, 92:1524

epidural sufentanil-ropivacaine, 92:180

levobupivacaine sparing by fentanyl, 92:410

preliminary investigation of remifentanyl (letter), 92:1358

sufentanil for PCEA, 92:184

lack of, associated with microfabricated microneedles, 92:502

low back, patients with, analgesic response to percutaneous neuromodulation therapy, 92:483

neuropathic

hyperalgesia, glucocorticoid inhibition, 92:476

response to carbamazepine and sustained-release morphine, patients pretreated with spinal cord stimulation, 92:488

nociception

Ca⁺⁺ channel blockers and morphine, 92:239

inflammatory pain and acute tolerance to alfentanil, 92:1296

orchialgia, pelvic hematoma after ilioinguinal nerve block, 92:1048

patients with and without, effect of epidural block on lymphocyte subsets and natural-killer cell activity, 92:463

postoperative

after intrathecal neostigmine, effect of dexamethasone, 92:228

analgesic requirements, effect of preemptive use of

dextromethorphan, 92:739

arthroscopy, knee, preemptive administration of antiinflammatory irrigation solution, 92:1301

cesarean delivery, effect of intravenous ketorolac, 92:1010

colonic surgery, epidural infusions of ropivacaine, 92:1276

effects of small-dose ketamine on propofol sedation, 92:1465

hand surgery, ambulatory,

postoperative analgesia at home, controlled comparison of tramadol, metamizol, and paracetamol, 92:347

hyperalgesia, EP₁ antagonist, 92:233

intrathecal sufentanil analgesia, 92:244

knee surgery, arthroscopic, efficacy of intraarticular sufentanil administration, 92:625

knee surgery, arthroscopic, preemptive analgesic effect of intraarticular morphine, 92:923

knee surgery, dextromethorphan as adjunct to PCA with morphine, 92:448

ondansetron and tramadol consumption, 92:1319

severe, comparison of two constant-dose continuous infusions of remifentanyl, 92:715

spinal fusion surgery, effect of intravenous propacetamol on postoperative morphine consumption, 92:1473

survey of patients after PCA, 92:224

thoracic epidural anesthesia and breathing, 92:1015

thoracotomy, effects of intrathecal morphine, sufentanil, and their combination vs IV morphine PCA, 92:31

thyroid surgery, bilateral superficial cervical plexus blocks, 92:1538

- total hip arthroplasty, continuous vs PCA femoral nerve sheath block, 92:455
- total hip or knee replacement, propacetamol vs ketorolac, 92:1569
- total knee arthroplasty, epidural small dose S(+)-ketamine, 92:1290
- posttraumatic, prehospital situation, tramadol as alternative to morphine, 92:1543
- prolonged, tunneled epidural catheters, pediatric patients, 92:1432
- reflex sympathetic dystrophy, glucocorticoid inhibition of hyperalgesia, 92:476
- thermal
analgesic interaction between intrathecal clonidine and glutamate receptor antagonists, rats, 92:725
hyperalgesia, effect of $\alpha 1$ - and $\alpha 2$ -adrenergic antagonists, 92:1558
- Pain scores**, see Measurement techniques
- Paracetamol**, see Analgesics, acetaminophen
- Paradoxical fat embolism syndrome**, see Complications
- Paralysis**, see Spinal cord
- Paraplegia**, see Complications
- Parathyroid surgery**, see Surgery
- Paravertebral block**, see Anesthetic techniques
- Parkinson's disease**, see Complications
- Parkinson's nocturnal hemoglobinuria**, see Complications
- Patent foramen ovale**, see Complications
- Patient-controlled analgesia (PCA)**, see Anesthetic techniques
- Patient-controlled epidural analgesia (PCEA)**, see Anesthetic techniques
- Pediatric anesthesia**, see Anesthesia
- PEEP**, see Ventilation, positive end-expiratory pressure
- Pelvic hematoma**, see Complications
- Pentalyte**, see Fluid balance
- Peptides**, adrenomedullin, mature, plasma concentration, and jugular venous oxygen saturation, during and after CPB, 92:837
- Perception**, effects of small-dose ketamine on propofol sedation, 92:1465
- Percutaneous neuromodulation therapy**, see Anesthetic techniques, electroanalgesia
- Peripheral nerve block**, see Anesthetic techniques
- Peripheral neuropathy**, see Complications
- Peripheral venous pressure**, see Blood pressure
- Personality**, see Psychology
- Personnel**
occupational exposure to inhaled anesthetics, 92:118
PACU staffing analysis, 92:947
weekday OR staffing analysis, 92:1493
- Pharmacodynamics**, heparin, onset time, 92:1391
- Pharmacokinetics**
adrenaline, endotracheal, beta adrenergic effect, 92:1408
aprotinin, plasma concentration, dose management, 92:830
gender and ϵ -aminocaproic acid, 92:1384
midazolam, oral, effect of anxiety and personality, 92:621
plasma concentration, lidocaine, effects of epidural insertion site and surgical procedure, 92:470
- Pharmacology**
acadesine, protective effect on lung ischemia-reperfusion injury, 92:590
angiotensin II antagonists
discontinuation before surgery, 92:26
discontinuation before surgery, against (letter), 92:CM56
aprotinin, regional hemostatic status and blood requirements after total knee arthroplasty, 92:775
calcium channel blockers, N-type, morphine and, combined effects on A δ vs C fiber mediated nociception, 92:239
carbachol, administered into brain stem pontine reticular formation and subarachnoid space, antinociceptive and sedative effects, rats, 92:1307
carbamazepine, response of neuropathic pain and pain in complex regional pain syndrome I, patients pretreated with spinal cord stimulation, 92:488
clonidine
intrathecal, and glutamate receptor antagonists, analgesic interaction between, thermal and formalin-induced pain, rats, 92:725
treatment of postanesthetic shivering, 92:257
colforsin daropate, effect on contractility in fatigued canine diaphragm, 92:762
dexamethasone
effect on postoperative pain and emesis after intrathecal neostigmine, 92:228
for preventing PONV associated with epidural morphine, dose-ranging study, 92:745
in ambulatory anesthesia, 92:85
dextromethorphan
large-dose, as adjunct to PCA with morphine after knee surgery, 92:448
preemptive use, effect on postoperative analgesic requirements, 92:739
diazepam cross-linked hemoglobin effects on tone of human saphenous vein, 92:324
large-dose infusions, hemoglobin and methemoglobin concentrations after, 92:44
diltiazem, effects on hemodynamics and seizure duration during ECT, 92:1327
epinephrine
ropivacaine with, as epidural test dose in parturients, limitations, 92:1529
with or without lidocaine, scalp infiltration, cardiovascular responses, craniotomy, 92:1516
esmolol, as alternative to remifentanyl, during desflurane anesthesia for fast-track outpatient gynecologic laparoscopic surgery, 92:352
ethanol
effects on Ca²⁺ sensitivity, airway smooth muscle, 92:767
enhancement of functional recovery of stunned myocardium independent of K_{ATP} channels, dogs, 92:299
glucocorticoids, methylprednisolone, inhibition of neuropathic hyperalgesia and spinal Fos expression, 92:476
heparin
rapid clinical onset, 92:1391
response to, effect of low preoperative antithrombin activity, cardiac surgery patients, adult vs infants, 92:66
hyaluronidase, in ophthalmic anesthesia (letter), 92:560
indigo carmine
-induced bradycardia, during general anesthesia, 92:276
vaginal, -induced severe hypotension (letter), 92:556
insulin
cardioplegia, effect on post-CABG arrhythmias, 92:810
infusion, effect on bupivacaine cardiotoxicity (letter), 92:37
ketamine, large concentrations, *in vitro* effects, attributed to nonspecific cytostatic effect, 92:927
meperidine, treatment of postanesthetic shivering, 92:257

Pharmacology—continued

- midazolam, dose-response
 - characteristics for reducing diaphragmatic contractility, 92:1590
- milrinone, administration before aortic cross-clamping, and intramyocardial cAMP during CPB, 92:676
- neostigmine, intrathecal, postoperative pain and emesis after, effect of dexamethasone, 92:228
- ondansetron
 - impairment of analgesic efficacy of tramadol, 92:1319
 - vs supplemental intraoperative oxygen, for prevention of PONV, 92:112
- ONO-8711, peripheral administration, effects on postoperative pain, rat model, 92:233
- oxycodone, administered into brain stem pontine reticular formation and subarachnoid space, antinociceptive and sedative effects, rats, 92:1307
- phenolamine, relief of thermal hyperalgesia, experimental mononeuropathy from chronic constriction injury, 92:1558
- phosphodiesterase III inhibitors
 - administration before aortic cross-clamping, and intramyocardial cAMP during CPB, 92:676
- olprinone, effects on hepatic vascular bed, endotoxemia, porcine model, 92:676
- prazosin, relief of thermal hyperalgesia, experimental mononeuropathy from chronic constriction injury, 92:1558
- propofol, induction dose, determinants, 92:656
- prostaglandin E₁, continuous infusion, and epidural analgesia, patient with paroxysmal nocturnal hemoglobinuria (letter), 92:1618
- RSR13, allosteric modification of oxygen delivery by hemoglobin, 92:615
- sevoflurane, patients with renal failure, 92:650
- SKF86466, relief of thermal hyperalgesia, experimental mononeuropathy from chronic constriction injury, 92:1558
- tacrolimus (FK506 506), effects on neurologic and histopathologic outcome after transient spinal cord ischemia, rats, 92:1237
- thiopental, inhibition of human $\alpha 7$ nicotinic acetylcholine receptor, 92:930
- tranexamic acid

- plasma concentrations during CPB, 92:1131
- regional hemostatic status and blood requirements after total knee arthroplasty, 92:775
- vasopressin, exogenous vs endogenous, during experimental CPR, 92:1499

Phentolamine, see Pharmacology**Phosphatidylinositol response, see**

Trachea

Phosphodiesterase III inhibitors, see Pharmacology**Platelet function, see Coagulation****Plethysmography, see Monitoring****Pneumocephalus, see Complications****Pneumonia, see Complications****Polycythemia rubra vera, see Complications****Popliteal fossa, see Anatomy****Position**

- lithotomy, knees-to-abdomen "Heimlich maneuver," morbidly obese patient (letter), 92:1619
- prone, effects on intraabdominal pressure and cardiovascular and renal function, patients with acute lung injury, 92:1226
- sitting, neuroanesthesia, monitoring, Germany, 1997 vs 1991, 92:166

Positive end-expiratory pressure, see Ventilation**Postanesthesia care unit (PACU)**

- air quality, online monitoring by proton-transfer-reaction mass spectrometry, 92:389
- computerized anesthesia records, score evaluation, 92:1203
- discharge from, administrative and medical delays, impact on total patient care hours, statistical analysis, 92:1222
- staffing, statistical analysis, 92:947

Postdural puncture headache, see Complications**Potassium, see Channels****Potency, anesthetic**

- minimum alveolar concentration (MAC)
 - enflurane, for LMA extubation, deeply anesthetized children, 92:72
 - for airway occlusion, 92:749
- hydrocarbons, effect of rigidity, shape, unsaturation, and length, 92:1477
- isoflurane, immobilizing action, modulation by cerebral and spinal GABA_A receptors, 92:1585
- isoflurane, mediators, GABA_A and strychnine-sensitive glycine receptors, 92:123
- volume local anesthetics, 92:205

Power analysis, see Statistics**Prazosin, see Pharmacology****Preeclampsia, see Complications****Preemptive analgesia, see Anesthetic techniques****Premedication**

- clonidine, oral, vs oral midazolam, pediatric tonsillectomy patients, 92:56
- midazolam
 - oral, vs oral clonidine, pediatric tonsillectomy patients, 92:56
 - rectal, vs rectal S(+)-ketamine, children, 92:62

Pressure

- intraabdominal, effects of prone positioning, patients with acute lung injury, 92:1226
- transdiaphragmatic, dose-response characteristics of midazolam, 92:1590

Pressure bags, see Equipment**Profound hypothermic circulatory arrest, see Hypothermia****Prone position, see Position****Propacetamol, see Analgesics, acetaminophen, prodrug****Propofol, see Anesthetics, intravenous****Prostaglandin E₁, see Pharmacology****Prostaglandin E receptors, see Receptors****Protamine, see Coagulation****Proteins**

- c-Fos, expression, ketamine-induced, in posterior cingulate and retrosplenial cortices, effects on xenon and N₂O, rats, 92:362
- coagulation protein function, effect of NO, thromboelastographic assessment, rabbits, 92:320
- heat shock protein 70, induction, time course, after brief period of ischemia in spinal cord, rabbits, 92:418

Proton-transfer-reaction mass**spectrometry, see Measurement techniques****Psychology**

- anxiety
 - effect on pharmacokinetics of oral midazolam, 92:621
- sensory stimuli and, children undergoing surgery, 92:897
- treatment, potential, auricular acupuncture, 92:548
- personality, effect on pharmacokinetics of oral midazolam, 92:621

Publications

- Anesthesia & Analgesia*, Internet technology and innovation (editorial), 92:1363

- journals, systematic reviews, critical appraisal, 92:700
- Pulmonary arterial pressure**, see Monitoring
- Pulmonary artery catheter**, see Equipment, catheters
- Pulmonary edema**, see Lung
- Pyridostigmine**, see Antagonists, neuromuscular relaxants
- Radiofrequency transmission**, see Electricity
- Rapacuronium**, see Neuromuscular relaxants
- Rapid sequence induction**, see Anesthetic techniques
- Recall**, intraoperative
explicit intraoperative recall at BIS of 47, 92:808
search for reliable awareness monitor (editorial), 92:801
- Receptors**
 α_1 -adrenergic, antagonists, relief of thermal hyperalgesia, experimental mononeuropathy from chronic constriction injury, 92:1558
 α_2 -adrenergic, agonists
clonidine, and glutamate antagonists, 92:725
clonidine, combined with bupivacaine, brachial plexus block, postoperative analgesia and hemodynamic changes, 92:199
clonidine, oral, vs oral midazolam, pediatric tonsillectomy patients, 92:56
clonidine, spinal, and lidocaine, antinociceptive interaction between, rat formalin test, isobolographic analysis, 92:733
 α_2 -adrenergic, antagonists, relief of thermal hyperalgesia, experimental mononeuropathy from chronic constriction injury, 92:1558
 β -adrenergic, agonists, endotracheal adrenaline, 92:1408
GABA_A
blockade, local, reversal of suppressive effects of isoflurane on thalamic neurons *in vivo*, 92:1578
cerebral and spinal, modulation of immobilizing capacity of isoflurane, 92:1585
mediation of MAC for isoflurane, 92:123
glutamate, antagonists, and intrathecal clonidine, analgesic interaction between, thermal and formalin-induced pain, rats, 92:725
glycine, strychnine-sensitive, mediation of MAC for isoflurane, 92:123
 μ -opioid, Ca²⁺ channel blockers and morphine, 92:239
muscarinic
dose-response study of anticholinesterase drugs, rats, 92:100
supraspinal and spinal action of cholinomimetics, 92:1307
nicotinic
acetylcholine, inhibition by thiopental, 92:930
supraspinal and spinal action of cholinomimetics, 92:1307
NMDA
antagonists, analgesic potency, question of mechanism-based use and timing (letter), 92:795
antagonists, and adenosine triphosphate, anesthetic interaction, rats, 92:134
antagonists, clonidine and glutamate antagonists, 92:725
antagonists, dextromethorphan, for pain relief after knee surgery, 92:448
antagonists, dextromethorphan, preemptive use, effect on postoperative analgesic requirements, 92:739
antagonists, xenon and N₂O, effects on ketamine-induced c-Fos expression in posterior cingulate and retrosplenial cortices, rats, 92:362
function, modulation by ketamine and magnesium, 92:1173
function, modulation by ketamine and magnesium, interactions with volatile anesthetics, 92:1182
opioid
inhibition of Ca²⁺ transients by morphine, 92:602
supraspinal and spinal action of cholinomimetics, 92:1307
prostaglandin E, subtype EP₁, peripheral administration, effects on postoperative pain, rat model, 92:233
- Recovery**
early discharge, considerations, awake craniotomy for removal of intracranial tumor, 92:89
facilitation, after day-case knee arthroscopy, and intraarticular sufentanil administration, 92:625
neurological, blood replacement with varying hematocrit levels, porcine model of profound hypothermic circulatory arrest, 92:329
- profiles, after sevoflurane anesthesia with or without intranasal fentanyl, pediatric patients, 92:1164
propofol vs midazolam, mechanically ventilated critically ill patients, 92:975
quality of, after anesthesia and surgery, scoring systems (letter), 92:281
time to discharge, after anorectal surgery, effect of dexamethasone, 92:85
- Rectal premedication**, see Anesthetic techniques
- Recurrent laryngeal nerve**, see Monitoring
- Recurrent laryngeal nerve palsy**, see Complications
- Reflex sympathetic dystrophy**, see Pain
- Reflex tachycardia**, see Heart
- Regional anesthesia and pain management**, see Anesthesia
- Regurgitation**, tricuspid or pulmonic valve, and pulmonary artery catheters, 92:1117
- Remifentanyl**, see Analgesics
- Renal toxicity**, see Toxicity, nephrotoxicity
- Respiration**, spontaneous, effects of small-dose ketamine on propofol sedation, 92:1465
- Respiratory events**, see Complications
- Resuscitation**, see Cardiopulmonary resuscitation
- Retroperitoneal lymph node dissection**, see Surgery
- Risk**, major, abdominal operations, caudal anesthesia, infants, 92:1439
- Ropivacaine**, see Anesthetics, local
- RSR13**, see Pharmacology
- Safety**
electroshock hazard, electrocautery and monitoring devices, 92:384
intraoperative TEE, 92:1126
ketorolac and sevoflurane anesthesia, 92:1058
sevoflurane, infants and children with congenital heart disease, 92:1152
- Sameridine**, see Analgesics; Anesthetics, local
- Saphenous vein**, see Veins
- Scalp infiltration**, see Anesthetic techniques
- Sciatic nerve**, see Nerve
- Sciatic nerve block**, see Anesthetic techniques
- Sedation**
prolonged, with propofol, effects on sleep, rats, 92:1232
propofol, effects of small-dose ketamine, 92:1465

Sedation—continued

propofol vs midazolam, mechanically ventilated critically ill patients, 92:975

supraspinal and spinal action of cholinomimetics, 92:1307

Seizure duration, during ECT, effects of diltiazem, 92:1327

Sepsis, see Infection

Serotonin

activity, spinal cord, modulations, and development of morphine tolerance, 92:1563

area postrema, effects of propofol, rats, 92:934

Sevoflurane, see Anesthetics, volatile

Shivering, see Complications

Shock, hypovolemic, after radical surgery for esophageal cancer, prediction, 92:1146

Single-lung ventilation, see Ventilation

Sitting position, see Position

SKF86466, see Pharmacology

Skin, transdermal drug delivery, lack of pain associated with microfabricated microneedles, 92:502

Sleep, deprivation, and prolonged sedation with propofol, rats, 92:1232

Smoking, see Complications

Society for Intravenous Anesthesia (SIVA), name change (editorial), 92:1

Sodium, see Channels

Somatosensory evoked potentials, see Monitoring

Sonoclot analysis, see Coagulation

Sonography, see Measurement techniques

Sore throat, see Complications

Soy, see Diet

Spinal anesthesia, see Anesthetic techniques

Spinal cord

clonidine and glutamate antagonists, 92:725

fentanyl/remifentanyl for SSEPs monitoring, 92:1510

ischemia

time course of acquisition of ischemic tolerance and induction of heat shock protein 70, rabbits, 92:418

transient, induced by aortic cross-clamping, effects of FK506 on neurologic and histopathologic outcome, rats, 92:1237

neurons, Fos expression, glucocorticoid inhibition, 92:476

paralysis, myelopathy and laminectomy, 92:1254

serotonin activity, modulations, and development of morphine tolerance, 92:1563

space deformation, laryngoscopy and, 92:1331

Spinal cord stimulation, see Anesthetic techniques

Spino-subcutaneous fistula, see Complications

Statistics

analysis of PACU staffing, 92:947

intravascular volume before obstetric spinal anesthesia, 92:997

median and minimum values, clarification (letter), 92:1073

meta-analysis

propofol vs midazolam in ICU, 92:975

systematic reviews in literature, 92:700

Monte-Carlo simulation, discharge from PACU, 92:1222

power analysis, more than two groups (letter), 92:797

weekday OR staffing analysis, 92:1493

Stereotactic frame placement, see Surgery

Stress response

immunosuppression early on during surgery, 92:1322

infants undergoing cardiac surgery, 92:882

multimodal postoperative rehabilitation program, 92:1594

Stroke volume variation, see Monitoring

Subarachnoid hemorrhage, see Brain

Subcutaneous infiltration, see Anesthetic techniques

Succinylcholine, see Neuromuscular relaxants

Sufentanil, see Analgesics

Superior laryngeal nerve injury, see Complications

Surgery

abdominal

aortic aneurysm repair, endovascular, patient undergoing, sudden respiratory arrest resulting from brain stem embolism, 92:335

infrarenal aneurysm repair, with large blood loss, successful use of hemoglobin-based oxygen carrier as primary blood substitute, 92:1423

major, caudal anesthesia, high-risk infants, 92:1439

major, intravascular volume replacement and coagulation, 92:565

major, postoperative pain and opioid requirements, bupivacaine wound instillation via PCA device and double-catheter system, 92:189

major, prolonged postoperative hypercoagulability, monitoring, thromboelastography, 92:572

repair of acute type A aortic dissection, hypoxemic respiratory failure after, successful treatment, 92:872

anorectal, outpatient, time to discharge, effect of dexamethasone, 92:85

brain, fluid responsiveness, stroke volume variation as predictor, 92:984

breast

ketorolac and sevoflurane anesthesia, 92:1058

modified radical mastectomy, continuous thoracic epidural vs general anesthesia, 92:1552

bronchoscopy, negative pressure post-tracheal extubation alveolar hemorrhage, 92:273

cardiac

congenital heart surgery, volume ventilation, comparison of two ventilators, infants, 92:76

remifentanyl (letter), 92:557

response to heparin, effect of low preoperative antithrombin activity, adults vs infants, 92:66

stress response, infants, 92:882

cardiopulmonary bypass (CPB)

adrenomedullin and jugular venous oxygen saturation, 92:837

differential perfusion, TEE imaging of new aortic cannula, 92:338

ϵ -ACA concentrations, lack of effect of gender, 92:1384

hypothermic, cognitive decline after, and high jugular bulb venous oxygen saturation, 92:1370

immediate or early extubation (letter), 92:1073

intramyocardial cAMP during, and inhibition of phosphodiesterase type III before aortic cross-clamping, 92:1377

mild hypothermic, jejunal mucosal perfusion, 92:5

plasma tranexamic acid concentrations, 92:1131

weaning from, difficult, prediction, diastolic dysfunction, 92:291

cervical laminectomy, posterior, acute transverse myelopathy after, 92:1254

cesarean delivery

effects of increase of central blood volume before spinal anesthesia, 92:997

intensity of labor pain, 92:1524

- parturient with Larsen syndrome, anesthetic management, 92:1266
- patient with polycythemia rubra vera and preeclampsia, epidural anesthesia, 92:1535
- postoperative pain and opioid requirement, effect of intravenous ketorolac, 92:1010
- cholecystectomy, diaphragm movement before and after, sonographic study, 92:755
- cleft palate repair, elective, postoperative opioid requirements, and prophylactically-administered rectal acetaminophen, 92:907
- colonic
 - colonoscopy, remifentanyl for monitored anesthesia care, 92:80
 - prolonged epidural infusions of ropivacaine after, impact of adding fentanyl, 92:1276
- coronary artery bypass graft (CABG)
 - apolipoprotein E polymorphisms and age at first CABG, 92:824
- arrhythmias, postoperative, effect of insulin, 92:810
- early extubation, remifentanyl vs fentanyl when combined with isoflurane/propofol, 92:1084
- fast-track, remifentanyl and fentanyl, efficacy and resource utilization, 92:1094
- fast-tracking after (editorial), 92:1081
- heart failure, postoperative, predicting risk of death, 92:596
- immediate or early extubation (letter), 92:1073
- craniotomy
 - awake, for removal of tumor, 92:89
 - awake, use of dexmedetomidine infusion, 92:1251
 - elective, peripheral venous pressure as hemodynamic variable, 92:172
 - hemodynamics during scalp infiltration, 92:1516
- esophagectomy, single-lung ventilation, intraoperative monitoring of recurrent laryngeal nerve, 92:662
- extracorporeal shock wave lithotripsy, nefopam analgesia as substitute for neurolept analgesia (letter), 92:557
- gynecologic
 - laparoscopic, fast-track outpatient, esmolol as alternative to remifentanyl during desflurane anesthesia, 92:352
 - PONV, ondansetron vs supplemental intraoperative oxygen, 92:112
- hand, ambulatory, postoperative analgesia at home, comparison of tramadol, metamizol, and paracetamol, 92:347
- hepatectomy
 - effects on plasma lidocaine concentration, 92:470
 - lactate and indocyanine green elimination rate in cirrhosis, 92:1064
- hysterectomy, dexamethasone and emesis, 92:745
- knee
 - arthroplasty, inhibition of synovial plasma extravasation, 92:1301
 - arthroplasty, total, postoperative pain, reduction, small-dose S(+)-ketamine applied with ropivacaine in epidural anesthesia, 92:1290
 - arthroplasty, total, regional hemostatic status and blood requirements with and without tranexamic acid or aprotinin, 92:775
 - arthroplasty, total, treatment of postoperative pain, propacetamol vs ketorolac, 92:1569
 - arthroscopic, ambulatory, preemptive analgesic effect of intraarticular bupivacaine and morphine, 92:923
 - arthroscopy, ambulatory anesthesia (letter), 92:556
 - arthroscopy, day-case, recovery after, and intraarticular sufentanil administration, 92:625
 - postoperative pain, dextromethorphan as adjunct to PCA with morphine, 92:448
- laparoscopic
 - fast-track outpatient gynecologic, esmolol as alternative to remifentanyl during desflurane anesthesia, 92:352
 - PONV, preventing, use of transcutaneous acupoint electrical stimulation, 92:629
- laparotomy
 - PCA wound instillation, 92:189
 - wound infiltration, local anesthetics (editorial), 92:3
- LASIK procedures, anesthetic interference with laser function, children, 92:1444
- liver transplantation, orthotopic antifibrinolytic therapy and pulmonary thromboembolism (letter), 92:559
- aprotinin administration and pulmonary thromboembolism, 92:1418
- mammoplasty, bilateral reduction, thoracic epidural anesthesia, patient with Klippel-Feil syndrome, 92:514
- ophthalmic
 - excimer laser procedures, anesthetic interference with laser function, children, 92:1444
 - hyaluronidase, in ophthalmic anesthesia (letter), 92:560
- orthopedic, major, coagulation, effects of Voluven vs HAES-steril, 92:855
- parathyroid, intravenous ketoprofen, 92:1052
- retroperitoneal lymph node dissection, for testicular cancer, anesthetic concerns (letter), 92:1618
- shoulder, major open, PCIA with ropivacaine 0.2% vs bupivacaine 0.15%, effects on hand motor function, 92:218
- spine
 - complex, peripheral venous pressure as hemodynamic variable, 92:172
 - fentanyl/remifentanyl for SSEPs monitoring, 92:1510
 - spinal fusion, postoperative morphine consumption, effect of intravenous administration of propacetamol, 92:1473
- stereotactic frame placement, nerve blocks vs subcutaneous infiltration, 92:424
- thoracic
 - anesthesia, pediatric, 92:1449
 - oxygenation during one-lung-ventilation, effects of inhaled NO and increasing levels of inspired fraction of oxygen, 92:842
 - preferred anesthetic technique, thoracic epidural anesthesia combined with general anesthesia, 92:848
- thoracoscopic mediastinal tumor resection, followed by reexpansion pulmonary edema, 92:1416
- thoracotomy, postoperative pain, effects of intrathecal morphine, sufentanil, and their combination vs IV morphine PCA, 92:31
- thyroid
 - intravenous ketoprofen, 92:1052
 - postoperative pain, prevention, bilateral superficial cervical plexus blocks, 92:1538
- tonsillectomy
 - PONV, effect of dexamethasone, 92:636

Surgery—continued

- postoperative coma, child with carnitine palmitoyltransferase I deficiency, 92:646
- preanesthetic medications, clonidine vs midazolam, 92:56
- total hip arthroplasty
 - embolic events during, 92:49
 - postoperative pain, relief, continuous vs PCA femoral nerve sheath block, 92:455
 - postoperative pain, treatment, propacetamol vs ketorolac, 92:1569
- transurethral prostatectomy, small-dose tetracaine spinal anesthesia, 92:1020
- urologic, multimodal postoperative rehabilitation program, 92:1594
- vascular
 - angiotensin II antagonists and anesthetic induction, 92:26
 - major, use of lepirudin for anticoagulation, patients with heparin-induced thrombocytopenia, 92:344

Surveys

- monitoring in Germany, 1997 vs 1991, 92:166
- patients, after PCA, 92:224

Sympathetic nervous system,

- adrenomedullin, mature, plasma concentration, and jugular venous oxygen saturation, during and after CPB, 92:837

Systemic lupus erythematosus, see

- Complications

Tacrolimus, see Pharmacology**Temperature**

- hypothermia
 - forced air warming vs negative pressure rewarming, hypothermic postanesthetic volunteers, 92:261
 - hypothermic CPB, cognitive decline after, and high jugular bulb venous oxygen saturation, 92:1370
 - mild hypothermic CPB, jejunal mucosal perfusion, 92:5
 - profound hypothermic circulatory arrest, porcine model, neurological recovery, blood replacement with varying hematocrit levels, 92:329
 - TEE detection of paradoxical fat embolism, 92:688
 - malignant hyperthermia, rapacuronium and ECT, 92:1171

Temporomandibular joint, see

- Complications

Thermoregulation

- forced air warming vs negative pressure rewarming, hypothermic postanesthetic volunteers, 92:261
- postanesthetic, shivering, treatment, comparison between meperidine, clonidine, and urapidil, 92:257

Thoracic epidural, see Anesthetic techniques**Thoracic surgery, see Surgery****Thoracoscopic mediastinal tumor**

- resection, see Surgery

Thoracotomy, see Surgery**Three vital capacity breaths, see**

- Anesthetic techniques

Thrombocytopenia, see Complications**Thromboelastography, see Coagulation;**

- Measurement techniques

Thromboembolism, see Complications**Thyroid surgery, see Surgery****Tolerance**

- acute, to alfentanil, and inflammatory pain, 92:1296
- morphine, development, and modulations of spinal serotonin activity, 92:1563

Tonsillectomy, see Surgery**Topical anesthesia, see Anesthetic techniques****Total hip arthroplasty, see Surgery****Tourniquet, see Equipment****Toxicity**

- cardiotoxicity
 - bupivacaine, favorable effect of insulin infusion (letter), 92:37
 - comparative, of S(-) and R(+) bupivacaine, and enantiomer-selective inhibition of L-type Ca^{2+} channels, 92:496
 - local anesthetic, resuscitation, 92:37
- cytotoxicity, cytostatic effect of ketamine, 92:927
- genotoxicity, waste anesthetic gases, 92:118
- nephrotoxicity
 - low-flow sevoflurane and isoflurane anesthesia, patients with stable moderate renal insufficiency, 92:650
 - safety of ketorolac and sevoflurane anesthesia, patients undergoing breast surgery, 92:1058
- neurotoxicity, lidocaine, obstetric patients (editorial), 92:287

Trachea, phosphatidylinositol response,

- dose-response study of anticholinesterase drugs, rats, 92:100

Tramadol, see Analgesics**Tranexamic acid, see Pharmacology****Transcranial Doppler ultrasonography, see Measurement techniques****Transcutaneous acupoint electrical**

- stimulation, see Antiemetic techniques

Transcutaneous CO₂, see Monitoring**Transdermal drug delivery, see Anesthetic techniques****Transesophageal echocardiography, see Monitoring****Transfusion, see Blood****Transient neurologic symptoms, see Complications****Transurethral prostatectomy, see Surgery****Trauma, see Brain****Tunneled epidural catheter, see Anesthetic techniques****Ultrasonography, see Imaging techniques****Unlighted stylet, see Equipment****Urapidil, see Pharmacology****Urologic surgery, see Surgery****Uterus, contractility, effects of opioids, 92:1006****Vascular endothelial growth factor, gene therapy with, for inoperable coronary artery disease, anesthetic management and results, 92:19****Vascular surgery, see Surgery****Vasodilation, endothelium-dependent, preservation, selective iNOS inhibition, sepsis, rats, 92:681****Vasopressin, see Pharmacology****Vecuronium, see Neuromuscular relaxants****Veins, saphenous, human, tone, effects of diaspirin cross-linked hemoglobin, 92:324****Ventilation**

- airway pressure release ventilation (APRV), gas exchange during, and CPAP, 92:950
- compromised, from TEE, proper probe positioning, infants (letter), 92:1076
- continuous positive airway pressure (CPAP)
 - effect on cerebral blood flow velocity, awake volunteers, 92:415
 - gas exchange during, and APRV, 92:950
- depression, remifentanyl infusion for severe postoperative pain, 92:715
- diaphragm
 - contractility, dose-response characteristics of midazolam, 92:762
- dysfunction, sepsis-induced, effects of ONO1714, hamsters, 92:959
- fatigued, contractility, effect of colforsin daropate, dogs, 92:762
- dyspnea, epidural anesthesia and respiratory distress, 92:749

- hypercarbic ventilatory response, effects of sameridine compared to morphine and placebo, 92:529
- mechanical
- acute lung injury and prone positioning, 92:1226
 - adhesion molecules and TGF- β 1 in ventilator-induced lung injury, 92:428
 - CO₂ monitoring, end-tidal vs transcutaneous, older children, 92:1427
 - comparison of two ventilators, infants, 92:76
 - constrictive pericarditis, 92:1424
 - critically ill patients, propofol vs midazolam, 92:975
 - stroke volume variation, fluids, and craniotomy, 92:984
- mechanics, effects of thoracic epidural analgesia with bupivacaine 0.25%, patients with severe chronic obstructive pulmonary disease, 92:1015
- one-lung, thoracic surgery
- oxygenation during, effects of inhaled NO and increasing levels of inspired fraction of oxygen, 92:842
 - oxygenation during, thoracic epidural anesthesia combined with general anesthesia, 92:848
 - pediatric patients, 92:1449
- positive end-expiratory pressure (PEEP), during active compression-decompression CPR with inspiratory threshold valve, 92:967
- pressure support, effect of breath termination criterion on breathing patterns and work of breathing, 92:161
- respiratory burst, lysophosphatidic acid and neutrophils, 92:1041
- single-lung, in esophagectomy, intraoperative monitoring of recurrent laryngeal nerve, 92:662
- transtracheal jet, and new intravenous catheter (letter), 92:1074
- work of breathing, effect of breath termination criterion during pressure support ventilation, 92:161
- Ventilators**, see Equipment
- Ventricular-peritoneal shunt**, see Equipment
- Vertebral metastasis**, see Complications
- Video imaging**, see Imaging techniques
- Vomiting, nausea and, postoperative (PONV)**
- after intrathecal neostigmine, effect of dexamethasone, 92:228
 - ambulatory surgery, oral ibuprofen vs codeine/acetaminophen, 92:1470
 - antiemetic effect of propofol, mechanism, 92:934
 - associated with epidural morphine, effect of dexamethasone, dose-ranging study, 92:745
 - intractable, after placement of permanent intrathecal catheter, treatment with caudal epidural blood patch, 92:1169
 - mastectomy, continuous thoracic epidural vs general anesthesia, 92:1552
 - patients' willingness to pay for antiemetic, 92:393
 - prediction of antiemetic rescue treatment, use of anesthetic information management system, 92:1203
 - prevention
 - after laparoscopic surgery, use of transcutaneous acupoint electrical stimulation, 92:629
 - ondansetron vs supplemental intraoperative oxygen, 92:112
 - use of dexamethasone, dose-ranging study, 92:745
- prophylaxis
- ondansetron and tramadol consumption, 92:1319
 - propofol vs desflurane-antiemetics, office-based anesthesia, 92:95
- thyroid and parathyroid surgery, intravenous ketoprofen, 92:1052
- tonsillectomy, effect of dexamethasone, 92:636
- Warming**, forced air, vs negative pressure rewarming, hypothermic postanesthetic volunteers, 92:261
- Weight**
- body size, effects on carbon monoxide concentrations in breathing circuit during low-flow anesthesia, 92:542
 - calculating induction dose of propofol, 92:656
- Weight-related dose regimen**, see Anesthetic techniques
- Western blot analysis**, see Measurement techniques
- Work of breathing**, see Ventilation
- World Federation of Societies of Anesthesiologists**, history, 92:1349
- Wound infiltration**, see Anesthetic techniques
- Xenon**, see Anesthetics, gases